

A Structure of Galois Extensions with an Inner Galois Group

George Szeto

Department of Mathematics
Bradley University
Peoria, Illinois 61625, USA
szeto@bradley.edu

Lianyong Xue

Department of Mathematics
Bradley University
Peoria, Illinois 61625, USA
lxue@bradley.edu

Abstract

Let B be a Galois extension of B^G with an inner Galois group G where $G = \{g_i \mid g_i(x) = U_i x U_i^{-1} \text{ for some } U_i \in B \text{ and for all } x \in B, i = 1, 2, \dots, n \text{ for some integer } n \text{ invertible in } B\}$. Then B is a composition of two Galois extensions $B \supset B^K$ with an inner Abelian Galois group K and $B^K \supset B^G$ with an inner Galois group G/K where $K = \{g \in G \mid g(U_i) = U_i \text{ for each } i\}$. Descriptions of $B \supset B^K$ and $B^K \supset B^G$ are given.

Mathematics Subject Classification: 13B05, 16W20

Keywords: Galois extensions, Galois algebras, central Galois algebras, Azumaya algebras, projective group algebras

1 Introduction

The class of central Galois algebras over a commutative ring with an inner Galois Group has been investigated ([1], [2]). It was shown that any central Galois algebra B over R with an inner Galois group G is an Azumaya projective group algebra RG_f of G over R with a factor set $f: G \times G \rightarrow \text{units of } R$ ([1], Theorem 6). Conversely, any Azumaya projective group algebra RG_f over

R with a factor set $f: G \times G \longrightarrow \text{units of } R$ is a central Galois algebra with an inner Galois group induced by the basis elements of RG_f ([2], Theorem 3). In case R is indecomposable, in [2], all central Galois algebras with an Abelian inner Galois group G are found, and an application to the commutative theory is obtained by giving a Kummer type theorem for Abelian extensions. Moreover some types of Galois extensions of a ring (not necessarily commutative) with an inner Galois group have been studied such as crossed products, skew polynomial rings, and Hirata separable extensions ([5], [8], [9], [10]). The purpose of the present paper is to continue the study of a Galois extension B of a ring B^G with an inner Galois group G where $G = \{g_i \mid g_i(x) = U_i x U_i^{-1} \text{ for some } U_i \in B \text{ and for all } x \in B, i = 1, 2, \dots, n \text{ for some integer } n \text{ invertible in } B\}$. We shall show that B is a composition of two Galois extensions: (1) $B \supset B^K$ with an Abelian inner Galois group K as studied in [2], where $K = \{g \in G \mid g(U_i) = U_i \text{ for each } i\}$ such that B^K is a central algebra over CK_f and contains B^G and CG_f as commutator subalgebras of B over C , and (2) $B^K \supset B^G$ with an inner Galois group G/K . Moreover, characterizations are given for B^K being a projective group ring of G/K over B^G .

Throughout this paper, we assume that B is a ring with 1, G a finite automorphism group of B of order n for some integer n invertible in B , C the center of B , B^G the set of elements in B fixed under each element in G . We call B a Galois extension of B^G with Galois group G if there exist elements $\{a_i, b_i \text{ in } B, i = 1, 2, \dots, m\}$ for some integer m such that $\sum_{i=1}^m a_i g(b_i) = \delta_{1,g}$ for each $g \in G$. A ring B is called a Galois algebra over R if B is a Galois extension of R which is contained in C , and a central Galois algebra over C if B is a Galois extension of C . Let A be a subring of B with the same identity 1. The commutator subring of A in B is denoted by $V_B(A)$. For the definitions of a separable or a Hirata separable extension and an Azumaya algebra, see [3] and [7]. As given in [1], a projective group ring RG_f of a group G of order n over a ring R with a factor set $f: G \times G \longrightarrow \text{units of } R$ is a ring with an R -basis $\{U_i, i = 1, 2, \dots, n \mid rU_i = U_i r \text{ for each } r \in R \text{ and } U_i U_j = U_k f(g_i, g_j) \text{ where } g_i g_j = g_k \in G\}$.

2 A Composition of Galois Extensions

In this section, let B be a Galois extension of B^G with an inner Galois group G of order n invertible in B , C the center of B , and $K = \{g \in G \mid g(U_i) = U_i \text{ for each } i = 1, 2, \dots, n\}$. Then we shall show that B is a composition of two Galois extensions: (i) B is a Galois extension of B^K with an Abelian inner Galois group K such that B^K is a central algebra over CK_f where CK_f is a projective group algebra of K over C with a factor set $f: K \times K \longrightarrow \text{units of } C$, and B^K contains B^G and CG_f as commutator subalgebras of B over C ,

and (ii) $B^K \supset B^G$ with an inner Galois group G/K . We begin with a lemma to show that B contains a projective group algebra CG_f , and that G contains a normal Abelian inner subgroup K .

Lemma 2.1 *Let B be a Galois extension of B^G with an inner Galois group G , $G = \{g_i \mid g_i(x) = U_i x U_i^{-1}, U_i \in B \text{ and for all } x \in B, i = 1, 2, \dots, n \text{ for some integer } n \text{ invertible in } B\}$. Then (i) B contains a separable projective group algebra CG_f of G over C with factor set $f : G \times G \rightarrow \text{units of } C$ where $f(g_i, g_j) = U_i U_j U_k^{-1}$ and $g_i g_j = g_k$, and (ii) K is an Abelian normal subgroup of G .*

Proof. (i) By Theorem 2.1 in [10], B contains a separable projective group algebra CG_f of G over C with factor set $f : G \times G \rightarrow \text{units of } C$ where $f(g_i, g_j) = U_i U_j U_k^{-1}$ and $g_i g_j = g_k$. Moreover, since the order n of G is invertible in B , CG_f is separable over C .

(ii) Since CG_f is invariant under the action of G and $K = \{g_j \in G \mid g_j(U_i) = U_i \text{ for each } i = 1, 2, \dots, n\}$, $U_j U_i = U_i U_j$ for each $g_j \in K$ and each $i = 1, 2, \dots, n$, that is, $g_j g_i = g_i g_j$ and $f(g_j, g_i) = f(g_i, g_j)$ for each $g_j \in K$ and for all $i = 1, 2, \dots, n$. Thus K is a normal Abelian inner subgroup of G .

Theorem 2.2 *Let B be a Galois extension of B^G with an inner Galois group G of order n invertible in B , and C the center of B . Then B is a composition of two Galois extensions: (1) B is a Galois extension of B^K with an Abelian inner Galois group K such that B^K is a central algebra over CK_f and contains B^G and CG_f as commutator subalgebras of B over C , and (2) B^K is a Galois extension of B^G with an inner Galois group G/K .*

Proof. (1) Since B is a Galois extension of B^G with an inner Galois group G , B is a Galois extension of B^K with an Abelian inner Galois group K by Lemma 2.1-(ii). Next, we claim that B^K is a central algebra over CK_f and contains B^G and CG_f as commutator subalgebras of B over C . In fact, since B is a Galois extension of B^G with an inner Galois group G , B is a Hirata separable extension of B^G ([7], Corollary 3). Since B is a Galois extension of B^G again, B is left or right B^G -finitely generated projective. Moreover, since n is invertible in B by hypothesis, B^G is a direct summand of B as a B^G -bimodule. Hence $V_B(V_B(B^G)) = B^G$ and $V_B(B^G)$ is a separable C -algebra ([6], Theorem 1) for B^G is separable over itself. Noting that CG_f is a separable C -subalgebra of $V_B(B^G)$, we have $CG_f = V_B(V_B(CG_f))$ ([6], Theorem 1). But $V_B(CG_f) = B^G$, so $V_B(B^G) = CG_f$. Thus B^G and CG_f are commutator subalgebras of B over C . Similarly, the pair B^K and CK_f are commutator subalgebras of B over C because B is a Galois extension of B^K with an inner Galois group K of an order invertible in B . Thus $V_B(B^K) = CK_f \subset B^K$, so

$V_{B^K}(B^K) = CK_f$, that is, CK_f is the center of B^K ; and so B^K is a central algebra over CK_f .

(2) Since n is a unit in B and K is a normal subgroup of G by Lemma 2.1, B^K is a Galois extension of B^G with inner Galois group G/K .

3 The Galois Subring B^K

In this section, B^K as given in Theorem 2.2 is described. Equivalent conditions are given for B^K which is a projective group ring of G/K over B^G . Thus more properties of B^K can be found. We shall keep all notations as given in section 2 for a Galois extension B of B^G with an inner Galois group G of order n invertible in B .

Theorem 3.1 *The following statements are equivalent:*

- (1) *The projective group algebra CG_f is a central Galois algebra over CK_f with Galois group G/K ;*
- (2) *CG_f is a central algebra over CK_f ;*
- (3) *B^G is a central algebra over CK_f ;*
- (4) *$B^K = B^G(G/K)_{\bar{f}}$ which is a projective group ring of G/K over B^G with factor set $\bar{f} : G/K \times G/K \rightarrow$ units of C where \bar{f} is induced by f ;*
- (5) *$B^K = \sum_{g \in G} B^G U_g$.*

Proof. (1) \implies (2) It is clear.

(2) \implies (1) Let $S = \{U_{\bar{g}} \mid \bar{g} \in G/K\}$ where $U_{\bar{g}} = U_g$ for an element $g \in G$. We claim that S are linearly independent over CK_f . In fact, let $\sum_{U_{\bar{g}} \in S} a_{\bar{g}} U_{\bar{g}} = 0$ for some $a_{\bar{g}} \in CK_f$. Then $a_{\bar{g}} = \sum_{h \in K} c_{g,h} U_h$ for some $c_{g,h} \in C$; and so $\sum_{U_{\bar{g}} \in S} \sum_{h \in K} c_{g,h} U_h U_{\bar{g}} = 0$, that is, $\sum_{U_{\bar{g}} \in S} \sum_{h \in K} c_{g,h} f(h, g) U_{hg} = 0$. Thus $c_{g,h} f(h, g) = 0$. Since $f(h, g)$ is a unit in C , $c_{g,h} = 0$ for each $U_{\bar{g}} \in S$ and $h \in K$. Therefore $a_{\bar{g}} = 0$ for each $U_{\bar{g}} \in S$. This implies that S are linearly independent over CK_f ; and so $CG_f = (CK_f)(G/K)_{\bar{f}}$, a projective group algebra G/K over CK_f with factor set $\bar{f} : G/K \times G/K \rightarrow$ units of C induced by $f : G \times G \rightarrow$ units of C . Noting that n , the order of G , is a unit in C , we conclude that $(CK_f)(G/K)_{\bar{f}}$ is a separable CK_f -algebra. Moreover, by hypothesis, CK_f is the center of CG_f , so CG_f is an Azumaya CK_f -algebra. Consequently, $CG_f (= (CK_f)(G/K)_{\bar{f}})$ is a central Galois algebra over CK_f with Galois group G/K ([2], Theorem 3).

(2) \iff (3) By Theorem 2.2, B^G and CG_f are commutator subalgebras of B , so it can be verified that B^G and CG_f have the same center.

(1) \implies (4) Since CG_f is a central Galois algebra over CK_f with Galois group G/K , $B^G(CG_f)$ is a Galois extension of B^G with Galois group G/K . On the other hand, B^K is also a Galois extension of B^G with Galois group

G/K such that $B^G(CG_f) \subset B^K$, so $B^G(CG_f) = B^K$ from the fact that the Galois group of $B^G(CG_f)$ is induced by and isomorphic with G/K . Therefore $B^K = B^G(CG_f) = B^G(CK_f)(G/K)_{\bar{f}} = B^G(G/K)_{\bar{f}}$.

(4) \implies (5) It is clear because $CK_f \subset B^G$.

(5) \implies (2) By hypothesis, $B^K = \sum_{g \in G} B^G U_g$, so $B^K = B^G G_f = B^G(CG_f)$. By Theorem 2.2, B^G and CG_f are commutator subalgebras of B , so that $B^K = B^G(CG_f)$ implies that B^K , B^G , and CG_f have the same center. But the center of B^K is CK_f by Theorem 2.2 again, so CK_f is the center of CG_f .

Theorem 3.1 implies some consequences in two special cases: (i) $K = \langle 1 \rangle$, and (ii) B^G is a separable CK_f -algebra. In case $K = \langle 1 \rangle$, Theorem 3.1 derives a result for a projective group ring, $B = B^G G_f$ which generalizes the structure theorem for an Azumaya projective group algebra as given in [2]. In case B^G is a separable CK_f -algebra, Theorem 3.1 gives a stronger structure theorem for B^K .

Corollary 3.2 *By keeping the notations of Theorem 3.1, the following statements are equivalent:*

- (1) CG_f is a central Galois algebra over CK_f with Galois group G/K and B^G is a separable CK_f -algebra;
- (2) CG_f is a central algebra over CK_f and B^G is a separable CK_f -algebra;
- (3) B^G is an Azumaya algebra over CK_f ;
- (4) $B^K = B^G(G/K)_{\bar{f}}$ as Azumaya algebras over CK_f ;
- (5) $B^K = \sum_{g \in G} B^G U_g$ as Azumaya algebras over CK_f .

We note that B is an Azumaya C -algebra if and only if B^K is a separable C -algebra. We shall show that B is a Hirata separable extension of CG_f and CK_f respectively.

Theorem 3.3 *Assume B is an Azumaya C -algebra. Then B is a Hirata separable extension of both CG_f and CK_f respectively such that both CG_f and CK_f are direct summands of B as a bimodule over themselves, respectively.*

Proof. Since n is a unit in B , CG_f and CK_f are separable C -subalgebras of B . By hypothesis, B is an Azumaya C -algebra, so B is projective over CG_f and CK_f respectively ([3], Proposition 2.3, page 48). Hence B is a Hirata separable extension of both CG_f and CK_f respectively ([4], Theorem 1). Moreover, both CG_f and CK_f are separable C -subalgebras of the Azumaya algebra B , so they are direct summands of B as a bimodule over themselves.

We conclude the present paper with two examples to demonstrate Galois extensions with an inner Galois group and an Abelian inner Galois group, respectively.

Example 1. Let R be a central Galois algebra with Galois group G and $B = \Delta(R, G, f)$ the crossed product of G over R with a trivial factor set f . Then (1) B is a Galois extension of $B^{\overline{G}}$ with an inner Galois group \overline{G} induced by G ;

(2) the center of $\Delta(R, G, f) = R_0 =$ the center of $R^G G_f$ where R_0 is the center of R ;

(3) B contains $R_0 G_f$, a projective group algebra of G over R_0 ;

(4) $R_0 G_f$ is not a central Galois algebra with Galois group \overline{G} induced by G where $\overline{G} = G/K$ and K is the center of G ;

(5) $B^K = R^G G_f$; and

(6) $B \supset B^K$ is a Galois extension with an Abelian Galois group K but $B^K \supset B^G$ is not a Galois extension with Galois group G/K .

Example 2. Let R be a commutative Galois algebra with a cyclic Galois group $\langle \rho \rangle$ of order n invertible in R , and $B = R[x, \rho] = R[X, \rho]/\langle X^n - a \rangle$ for some a invertible in R^ρ where $\langle X^n - a \rangle$ is the ideal of $R[X, \rho]$ generated by $X^n - a$. Then

(1) $B = R[x, \rho]$ is a separable skew polynomial ring over R with a basis $\{1, x, x^2, \dots, x^{n-1} \mid x^n = a\}$ where $xr = \rho(r)x$ for any $r \in R$;

(2) $B \cong R\langle \rho \rangle_f$, a crossed product of $\langle \rho \rangle$ over R with factor set $f : \langle \rho \rangle \times \langle \rho \rangle \rightarrow$ units of R by $f(\rho^i, \rho^j) = \begin{cases} 1 & \text{if } i + j < n \\ a & \text{if } i + j \geq n; \end{cases}$

(3) $B \supset B^\rho$ is a Galois extension with an Abelian inner Galois group $\langle \overline{\rho} \rangle$ induced by ρ ;

(4) the center of B is R^ρ ;

(5) B contains $R^\rho \langle \overline{\rho} \rangle_f$, a projective group algebra of $\langle \overline{\rho} \rangle$ over R^ρ with factor set f ;

(6) $R^\rho \langle \overline{\rho} \rangle_f$ is a commutative separable R^ρ -algebra, so $K = \langle 1 \rangle$; and hence $B^K = B$.

ACKNOWLEDGEMENTS. This paper was written under the support of a Caterpillar Fellowship at Bradley University. The authors would like to thank Caterpillar Inc. for the support.

References

- [1] F.R. DeMeyer, Some Notes on the General Galois Theory of Rings, *Osaka J. Math.*, **2** (1965), 117-127.
- [2] F. R. DeMeyer, Galois Theory in Separable Algebras over Commutative Rings, *Illinois J. Math.*, **10** (1966), 287-295.

- [3] F. R. DeMeyer and E. Ingraham, Separable algebras over commutative rings, Volume 181. Springer Verlag, Berlin, Heidelberg, New York, 1971.
- [4] S. Ikehata, Note on Azumaya Algebras and H -Separable Extensions, *Math. J. Okayama Univ.*, **23** (1981), 17-18.
- [5] S. Ikehata, On H -separable polynomials of prime degree, *Math. J. Okayama Univ.* **33** (1991), 21-26.
- [6] K. Sugano, On Centralizers In Separable Extensions II, *Osaka J. Math.*, **8** (1971), 465-469.
- [7] K. Sugano, On a Special Type of Galois Extensions, *Hokkaido J. Math.*, **9** (1980), 123-128.
- [8] G. Szeto and L. Xue, Skew Group Rings which are Galois, *International Journal of Mathematics and Mathematical Sciences*, **23**(4) (1999), 279-283.
- [9] G. Szeto and L. Xue, On Splitting Rings for Azumaya Skew Polynomial Rings, *Algebra Colloquium*, **8**(1) (2001), 25-32.
- [10] G. Szeto and L. Xue, On Galois Extensions with an Inner Galois Group, *Recent Developments in Algebra and Related Area*, ALM 8, 239-245, Higher Education Press and International Press Beijing-Boston, 2008.

Received: September, 2010