International Journal of Algebra, Vol. 7, 2013, no. 3, 131 - 144
HIKARI Ltd, www.m-hikari.com

On KUS-Algebras

Samy M. Mostafa[a], Mokhtar A. Abdel Naby[a], Fayza Abdel Halim[b]

and Areej T. Hameed[b]

[a] Department of Mathematics, Faculty of Education
Ain Shams University, Roxy, Cairo, Egypt

[b] Department of Pure Mathematics, Faculty of Sciences
Ain Shams University, Cairo, Egypt

samymostafa@yahoo.com, abdelnaby@hotmail.com, salahfi@hotmail.com
areej238@gmail.com

Abstract

The aim of this paper is to introduce and study new algebraic structure, called KUS-algebra and investigate some of its properties.

Mathematics Subject Classification: 06F35, 03G25

Keywords: KUS-algebra, KUS-sub-algebra, KUS-ideal, homomorphism of KUS-algebra, (p-semi-simple, medial) KUS-algebra

Copyright © 2013 S. M. Mostafa et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

The notion of BCK-algebras was proposed by Iami and Iseki in 1966. In the same year, K. Is´eki [4] introduced the notion of a BCI-algebra which is a generalization of a BCK-algebra. Since then numerous mathematical papers have been written investigating the algebraic properties of the BCK/BCI-algebras and their relationship

with other universal structures including lattices and Boolean algebras. There is a great deal of literature has been produced on the theory of BCK/BCI-algebras, in particular, emphasis seems to have been put on the ideal theory of BCK/BCI-algebras see[3]. For the general development of BCK/BCI-algebras the ideal theory plays an important role. Y. Komori ([6]) introduced a notion of BCC-algebras, and W. A. Dudek [1] redefined the notion of BCC-algebras by using a dual form of the ordinary definition in the sense of Y. Komori. In([1], [2]), C. Prabpayak and U. Leerawat introduced the concept of KU-algebra . They gave the concept of homomorphism of KU-algebras and investigated some related properties. In this paper the concepts of KUS-algebras, KUS-sub-algebras , KUS-ideals , homomorphism of KUS-algebras are introduced. The relation between some abelian groups and KUS-algebras , the G-part of KUS-algebras are studied and investigated some of its properties.

2. The Structure of KUS-algebras

In this section, we will introduce a new notion called KUS-algebras and study several properties of it.

Definition 2.1([1],[2]). A KU-algebra is a nonempty set X with a constant (0) and a binary operation $(*)$ satisfying the following axioms: for any $x , y , z \in X$,

(i) $(x*y) * [(y*z) * (x*z)]= 0$,
(ii) $0*x = x$,
(iii) $x*0 = 0$,
(iv) $x*y = 0$ and $y*x = 0$ imply $x = y$.

Lemma 2.2([1],[2]). Every KU-algebra X satisfies the following conditions:

(v) $x* (y*z) = y* (x*z)$, for arbitrary $x ,y ,z \in X$.
(vi) $x*x = 0$, for arbitrary $x \in X$.

Definition 2.3. Let $(X; *,0)$ be an algebra of type (2,0) with a single binary operation $(*)$. Then $(X; *,0)$ is called KUS-algebra if it satisfies the following axioms : for any $x , y , z \in X$,

$(kus_1) : (z*y) * (z*x) = (y*x)$,
$(kus_2) : 0*x = x$,
$(kus_3) : x*x = 0$,
$(kus_4) : x* (y*z) = y* (x*z)$.

In X we can define a binary relation $(\leq)$ by : $x \leq y$ if and only if $y * x = 0$.

A KU-algebra $(X; *,0)$ is called KUS-algebra if it satisfies:

(vii) $(z*y) * (z*x) = (y*x)$.

For brevity we shall call X a KUS-algebra unless otherwise specified

Example 2.4. Let X = {0, a, b, c, d} in which $(*)$ is defined by the following table:

$*$	**0**	**a**	**b**	**c**	**d**
0	**0**	**a**	**b**	**c**	**d**
a	**d**	**0**	**a**	**b**	**c**
b	**c**	**d**	**0**	**a**	**b**
c	**b**	**c**	**d**	**0**	**a**
d	**a**	**b**	**c**	**d**	**0**

It is easy to show that $(X; *,0)$ is KUS-algebra .

Now, we give some properties and theorems of KUS-algebras.

Proposition 2.5. Let X be a KUS-algebra. Then the following holds:
for any $x, y, z \in X$,

a) $x * y = 0$ and $y * x = 0$ imply $x = y$,
b) $x * (y * x) = y * 0$,
c) $(x * y) = 0$ implies that $x * 0 = y * 0$,
d) $(x*y) * 0 = y*x$,
e) $x*0 = 0$ implies that $x = 0$,
f) $x = 0 * (0 * x)$,
g) $0 * (x * y) = (0 * x) * (0 * y)$,
h) $x*z = y*z$ implies that $x*0 = y*0$.

Proof:

a) Since $x * y = 0$ and $y * x = 0$, then $x \leq y$ and $y \leq x$ imply $x = y$.
b) $x * (y*x) = y * (x*x) = y*0$.
c) $(0*y) * (x * y) = x * 0$, by (b), and $(0*y) * (x * y) = (0 * y) * 0$, since $(x * y) = 0$. Then $x * 0 = y * 0$.
d) $(x*y) * 0 = (x*y) * (x*x) = y*x$, by (kus_1).
(e), (f) and (g) are clears by (kus_2).
h) $x*0 = x* (z*z) = z* (x*z) = z*(y*z) = y*(z*z) = y*0$. ⌂

Proposition 2.6. Let X be a KUS-algebra . A relation $(\leq)$ on X defined by

$x \leq y$ if $y * x = 0$. Then $(X, \leq)$ is a partially ordered set.

Proof: Let X be a KUS-algebra and let x, y, z $\in$ X, since $x * x = 0$, $x \leq x$. Suppose that $x \leq y$ and $y \leq x$, then $x * y = 0 = y * x$. By proposition (2.5(a)), x = y. Suppose that $x \leq y$ and $y \leq z$, then $y * x = 0$ and $z*y = 0$. By (kus_1)
$0 = (y * x) * (y * x) = (y * x) * [(z * y) * (z * x)] = 0 * [0 * (z * x)] = z * x$, hence $x \leq z$. Thus $(X, \leq)$ is a partially ordered set. ⌂

Proposition 2.7. Let X be a KUS-algebra . Then the following holds: for any x, y, z $\in$X,

1. $x * y \leq z$ imply $z * y \leq x$,
2. $x \leq y$ implies that $z * x \leq z * y$,
3. $y * [(y * z) * z] = 0$,
4. $(x * z) * (y * z) \leq (y * x)$,
5. $x \leq y$ and $y \leq z$ imply $x \leq z$,
6. $x \leq y$ implies that $y * z \leq x * z$.

Proof:

1. It follows from (kus_4) .
2. By (kus_1), we obtain $[(z * y) * (z * x)] = (y * x)$, but $x \leq y$ implies $y * x = 0$, then we get $(z * y) * (z * x) = 0$. i.e., $z * x \leq z * y$.
3. It is clear by (kus_4) and (kus_3) .
4. By (kus_3) , (kus_4) and (kus_1) , $(y * x) * [(x * z) * (y * z)] = (x * z) * [(y * x) * (y * z)] = (x * z) * (x * z) = 0$. Thus $(x * z) * (y * z) \leq (y * x)$.
5. If $x \leq y$, then by (2), $z * x \leq z * y$. By applying (kus_2) and $(x \leq y)$, $z * x = 0 * (z * x) = (y * x) * (z * x) \leq z * y = 0$ [by (4) and $(y \leq z)$] imply $z * x \leq 0$, i.e., $0 * (z * x) = 0$. By (kus_2), $z * x = 0$ and so $x \leq z$.
6. If $x \leq y$, then $(x * z) * (y * z) = (y * x) = 0$. Hence $y * z \leq x * z$. ⌂

Proposition 2.8. Every KUS-algebra X satisfying $x * (x * y) = x * y$ for all x, y $\in$ X is a trivial algebra .

Proof: putting x = y in the equation $x * (x * y) = x * y$, we have $x*0 = 0$ .By (kus_2) , x = 0. Hence X is a trivial algebra .⌂

3. KUS-ideals and Homomorphism of KUS-algebras

In this section we will present some results on images and preimages of homomorphism on KUS-algebras.

Definition 3.1. Let X be a KUS-algebra and let S be a nonempty set of X. S is called a KUS-sub-algebra of X if $x * y \in S$ whenever $x, y \in S$.

Definition 3.2([5]). Let I be a nonempty subset of X , I is called an ideal of X if, for all $x, y \in X$

(I_1) $0 \in I$,

(I_2) $x \in I$ and $y * x \in I$ imply $y \in I$.

Definition 3.3([1],[2]). A nonempty subset I of a KU-algebra X is called a KU-ideal of X if it satisfies the following conditions : for all $x, y, z \in X$

(KU_1) $0 \in I$,

(KU_2) $x * (y * z) \in I$, $y \in I$ implies $(x * z) \in I$.

Definition 3.4. A nonempty subset I of a KUS-algebra X is called a KUS-ideal of X if it satisfies: for $x , y, z \in X$,

($Ikus_1$) $(0 \in I)$,

($Ikus_2$) $(z * y) \in I$ and $(y * x) \in I$ imply $(z * x) \in I$.

Example 3.5 . Let $X = \{0 ,a, b, c\}$ in which $(*)$ is defined by the following table:

*	**0**	**a**	**b**	**c**
0	**0**	**a**	**b**	**c**
a	**a**	**0**	**c**	**b**
b	**b**	**c**	**0**	**a**
c	**c**	**b**	**a**	**0**

Then $(X; *,0)$ is KUS-algebra . It is easy to show that $I_1 = \{0,a\}$, $I_2 = \{0,b\}$, $I_3 = \{0,c\}$, and $I_4 = \{0, a, b, c\}$ are KUS-ideals of X .

Proposition 3.6. Let X be a KUS-algebra and I be a nonempty subset of X containing **0**. Then I is a KUS-ideal of X if and only if :

$(z * y) \in I$, $(z * x) \notin I$ imply $(y * x) \notin I$, for all $x, y, z \in X$.

Proof: Let I be an KUS-ideal of X and $(z * y) \in I$, $(z * x) \notin I$. Suppose that $(y * x) \in I$, since I is a KUS-ideal, then $(z * x) \in I$, a contradiction .

Conversely, assume that $(z*y) \in I, (z*x) \notin I$ imply $(y*x) \notin I$, for all $x, y, z \in X$. If $(z*y) \in I, (y*x) \in I$. It is clear that $(z*x) \in I$. then I is a KUS-ideal of X. ⌂

Corollary 3.7.

Let X be a KUS-algebra and I be a nonempty subset of X containing **0**. Then :

(C_1) I is a KUS-ideal of X if and only if $(z*y) \in I, (y*x) \notin I$ imply $(z*x) \notin I$, for all $x, y, z \in X$.

(C_2) I is a KUS-ideal of X if and only if $(z*y) \in I, (z*x) \in I$ imply $(y*x) \in I$, for all $x, y, z \in X$.

(C_3) I is a KUS-ideal of X if and only if $(z*x) \in I, (y*x) \in I$ imply $(z*y) \in I$, for all $x, y, z \in X$.

Proposition 3.8. Every KUS-ideal of KUS-algebra X is a KUS-sub-algebra.

Proof: For all $x, y, z \in X$, let I be a KUS-ideal of a KUS-algebra X such that $x, y \in I$, then $(0*x) = x \in I, (0*y) = y \in I$. Hence, by corollary $(3.7(C_2))$, $x*y \in I$. Therefore I is a KUS-sub-algebra.⌂

Proposition 3. 9. Every KUS-ideal of X is an ideal of X.

Proof: For all $x, y, z \in X$, let I is KUS-ideal of X. By corollary$(3.7(C_3))$ $(z*x) \in I$ and $(y*x) \in I$ imply $(z*y) \in I$. If $z = 0$,then $(0*x) = x \in I$ and $(y*x) \in I$ imply that $(0*y) = y \in I$ **.** and hence I is an ideal of X. ⌂

In general, the converse of proposition (3.9) is not true . For example:

Example 3.10. Let $X = \{0,1, 2, 3\}$ be a KUS-algebra with the following Cayley table:

*	0	1	2	3
0	0	1	2	3
1	3	0	1	2
2	2	3	0	1
3	1	2	3	0

Then $(X; *, 0)$ is KUS-algebra . It is easy to show that $I = \{0, 3\}$ is an ideal of X which is not a KUS-ideal of X , since $3 * 2 = 3 \in I$, $2 * 1 = 3 \in I$ and $3 * 1 = 2 \notin I$.

Proposition 3.11. Every KUS-ideal of KUS-algebra X is a KU-ideal of X.

Proof: For all x, y, z ∈ X, let I is KUS-ideal of a KUS-algebra X. If x* (y* z)∈ I , y ∈ I, then y = 0∗ y = (z∗ z) ∗ y, by (kus_3).

= [y∗ (z∗ z)] ∗ 0 , by proposition(2.5(d)).

= [z∗ (y∗ z)] ∗ 0, by (kus_4).

= (y∗ z) ∗ z ∈ I , by proposition(2.5(d)).

Then x∗ (y∗ z)∈ I and (y∗ z) ∗ z ∈ I implies (x∗ z)∈ I .

Hence I is a KU-ideal of X. ⌂

In general, the converse of proposition (3.11) is not true .

For example: Let X ={0,1, 2, 3} be a KUS-algebra with the following Cayley table

∗	0	1	2	3
0	0	1	2	3
1	3	0	1	2
2	2	3	0	1
3	1	2	3	0

Then (X; ∗,0) is KUS-algebra . It is easy to show that I ={0, 3} is a KU-ideal of X ,which is not a KUS-ideal of X , since 3 ∗2 = 3 ∈ I , 2 ∗1 = 3 ∈I and 3∗ 1 = 2 ∉I.

Proposition 3.13. Let $\{I_i \mid i\in\Lambda\}$ be a family of KUS-ideals on KUS-algebra X. Then $\bigcap_{i\in\Lambda} I_i$ is a KUS-ideal of X.

Proof: Since $\{I_i \mid i\in\Lambda\}$ be a family of KUS-ideals of X , then $0 \in I_i$, for all i ∈Λ, then $0 \in \bigcap_{i\in\Lambda} I_i$. For any x, y, z ∈ X , suppose $z*y \in I_i$ and $y*x \in I_i$, for all i ∈Λ, but I_i is a KUS-ideal of X for all i ∈Λ. Then $z*x \in I_i$, for all i ∈Λ, therefore, $z*x \in \bigcap_{i\in\Lambda} I_i$. Hence $\bigcap_{i\in\Lambda} I_i$ is KUS-ideal of KUS-algebra X .⌂

Definition 3.14 ([2]). Let (X ; ∗,0) and (Y; ∗`,0`) be KUS-algebras , the mapping f : (X; ∗,0) → (Y; ∗`,0`) is called a homomorphism if it satisfies:

f (x∗ y) = f (x) ∗` f (y) for all x , y ∈ X.

Theorem 3.15. Let f : (X; ∗,0) → (Y; ∗`,0`) be into homomorphism of a KUS-algebras, then :

A) f (0) = 0'.

B) f is injective if and only if Ker $f = \{0\}$.

C) $x \leq y$ implies $f(x) \leq f(y)$.

Proof: Clear.

Theorem 3.16. Let $f : (X; *,0) \rightarrow (Y; *`,0`)$ be an into homomorphism of a KUS-algebras, then :

(F_1) If S is a KUS-sub-algebra of X, then f (S) is a KUS-sub-algebra of Y .

(F_2) If I is a KUS-ideal of X, then f (I) is a KUS-ideal in Y .

(F_3) If B is a KUS-sub-algebra of Y, then f^{-1}(B) is a KUS-sub-algebra of X .

(F_4) If J is a KUS- ideal in Y, then f^{-1} (J) is a KUS-ideal in X .

(F_5) Ker f is KUS-ideal of X.

(F_6) Im(f) is a KUS-sub-algebra of Y.

Proof: Clear .

4. The G-part of KUS-algebras

In this section we give some basic definitions , preliminaries and lemmas of G-part in KUS-algebras.

Definition 4.1. Let (X; $*$,0) be a KUS-algebra. For any nonempty subset S of X, we define $G(S) := \{x \in S \mid x * 0 = x\}$.

In particular, if S = X then we say that G(X) is the G-part of KUS-algebra X.

For any KUS-algebra X, the set $B(X) := \{x \in X \mid x * 0 = 0\}$ is called a p-radical of X.

A KUS-algebra X is said to be p-semi-simple if B(X) =$\{0\}$.

The following property is obvious: $G(X) \cap B(X) = \{0\}$.

Proposition 4.2. If (X; $*$,0) is a KUS-algebra and x, y$\in$X, then

$y \in G(X) \Leftrightarrow x * (y * x) = y$.

Proof: By (kus_4) and (kus_3) $x * (y * x) = y * (x * x) = y * 0 = y \Leftrightarrow y \in G(X)$.⌂

Corollary 4.3. If (X; $*$,0) is a KUS-algebra and x, y$\in$X, then y$\in$B(X)

$\Leftrightarrow x * (y * x) = 0$.

Proof: By (kus_4) and (kus_3) $x * (y * x) = y * (x * x) = y * 0 = 0 \Leftrightarrow y \in B(X)$.⌂

Proposition 4.4. If $(X; *,0)$ is a KUS-algebra and $b * a = c * a$, then $b * 0 = c * 0$, where $a, b, c \in X$.

Proof: By (kus_4) and (kus_3), $a * (b * a) = b * (a * a) = b * 0$ and $a * (c * a) = c * (a * a) = c * 0$. Since $b * a = c * a$, then $b * 0 = c * 0$.⌂

Corollary 4.5. Let X be a KUS-algebra. Then the right cancellation law holds in G(X).

Proof: Let $a, b, c \in G(X)$ with $b * a = c * a$. By proposition (4.4), $b * 0 = c * 0$. Since $b, c \in G(X)$, we obtain $b = c$.⌂

Proposition 4.6. Let $(X; *,0)$ be a KUS-algebra. Then
$x \in G(X)$ if and only if $x * 0 \in G(X)$.

Proof: If $x \in G(X)$, then $x * 0 = x$ and $(x * 0) * 0 = x * 0$. Hence $x*0 \in G(X)$. Conversely, if $x * 0 \in G(x)$, then $(x * 0) * 0 = x * 0$. By applying corollary (4.5), we obtain $x * 0 = x$. Therefore $x \in G(X)$.⌂

Corollary 4.7. Let X be a KUS-algebra. Then the left cancellation law holds in G(X).

Proof: Let $a, b, c \in G(X)$ with $a * b = a * c$. Then $a* y = a*(y*0) = y*(a*0) = y*a$, for any $y \in G(X)$. By proposition (4.6), $b = a * (b * a) = a * (a * b) = a * (a * c)$
$= a * (c * a) = c$, we obtain $b = c$.⌂

Proposition 4.8. Let $(X;*,0)$ be a KUS-algebra, for all $x, y, z \in G(X)$, then
$L_1)$ $y * x = z$ imply $x * z = y$,
$L_2)$ $x * (0 * y) = y * (0 * x)$.

Proof:
$L_1)$ Since $y * x = z$, then
$x * z = (y * x) * (y * z)$, by (kus_1).
$= z * (y * z)$, since $(y * x = z)$.
$= y * (z * z)$, by (kus_4).
$= y * 0 = y$, since $y \in G(X)$.
$L_2)$ Since $((y * (0 * x)) * x) * 0 = x * (y * (0 * x))$, by proposition(2.5(d)).
$= y * (x * (0 * x))$, by (kus_4).
$= y * (0 * 0)$, by proposition(2.5(b)).

$= y * 0 = y$, since $y \in G(X)$,

we have $((y * (0 * x)) * x) * 0 = y$. It follows from that $(y * (0 * x)) * x = 0 * y$ and hence $y * (0 * x) = x * (0 * y)$ by (L_1) . ⌂

The following theorem state the relation between KUS-algebra and abelian group

Theorem 4.9. Let $(X; \cdot, -1, e)$ be an abelian group. If $x * y = x^{-1} \cdot y$ is defined and $0 = e$, then $(X; *, 0)$ is a KUS-algebra .

Proof: We only show that conditions (kus_1) and (kus_4) of KUS-algebra are satisfied. For the case of (kus_1) , since X is an abelian group, we have

$(z * y) * (z * x) = (z^{-1} \cdot y)^{-1} \cdot (z^{-1} \cdot x) = (y^{-1} \cdot z) \cdot (z^{-1} \cdot x) = y^{-1} \cdot (z \cdot z^{-1}) \cdot x = y^{-1} \cdot x = (y * x)$

For the case of (kus_4) , we also have

$x * (y * z) = x^{-1} \cdot (y^{-1} \cdot z) = (x^{-1} \cdot y^{-1}) \cdot z = (y^{-1} \cdot x^{-1}) \cdot z = y^{-1} \cdot (x^{-1} \cdot z) = y * (x * z)$.⌂

Theorem 4.10. Let $(X; *, 0)$ be a KUS-algebra. If $x \cdot y = x * y$ is defined , $x^{-1} = x$ and $e = 0$, then the structure $(X ; \cdot , -1 , e)$ is an abelian group.

Proof: We only show that the structure $(X ; \cdot , -1 , e)$ satisfies the conditions of associative and commutative with respect to the operation $(\cdot)$.

For associative , we have

$x \cdot (y \cdot z) = x * (y * z)$

$= (0 * x) * (y * (0 * z))$, by (kus_2).

$= (0 * x) * (z * (0 * y))$, by proposition $(4.8(L_2))$.

$= z * ((0 * x) * (0 * y))$, by (kus_4).

$= z * (x * y)$, by (kus_2).

$= ((x * y) * z) * 0$, by proposition(2.5(d)).

$= (x * y) * z$

$= (x \cdot y) \cdot z$.

For commutative, we also have

$x \cdot y = x * y$

$= x * (0 * y)$

$= y * (0 * x)$, by proposition $(4.8(L_2))$.

$= y * x = y \cdot x$. ⌂

Definition 4.11. Let $(X; *, 0)$ be a KUS-algebra satisfying

$(x * y) * (z * u) = (x * z) * (y * u)$, for any x, y, z and $u \in X$ is called a medial of KUS-algebra.

Proposition 4.12. Let $(X; *,0)$ be a medial of KUS-algebra. Then G(X) is a KUS-sub-algebra of X .

Proof: Let $x, y \in G(X)$, then $x*0 = x$ and $y*0 = y$. Hence

$$\begin{aligned}(x*y)*0 &= (x*y) * (0*0)\\ &= (x*0) * (y*0)\\ &= x*y \ .\end{aligned}$$

⌂

Corollary 4.13. Let $(X; *,0)$ be a medial of KUS-algebra. Then B(X) is a KUS-sub-algebra of X .

Proof: Let $x, y \in B(X)$, then $x*0 = 0$ and $y*0 = 0$. Hence

$$(x*y)*0 = (x*y) * (0*0) = (x*0) * (y*0) = 0*0 = 0 \ .$$

⌂

Theorem 4.14. In a KUS-algebra $(X; *,0)$, X is medial of KUS-algebra if and only if it satisfies : for any $x, y, z \in G(X)$,

M_1) $(x*y) = (y*x)$,

M_2) $(x*y)*z = (x*z)*y$.

Proof: Suppose $(X; *,0)$ is medial of KUS-algebra and $x, y, z \in G(X)$. Then :

M_1) $y*x = 0*(y*x) = (x*x) * (y*x) = (x*y) * (x*x) = (x*y)*0 = (x*y)$.

M_2) $(x*y)*z = (x*y) * (z*0) = (x*z) * (y*0) = (x*z)*y$.

Conversely , assume that the conditions (M_1) and (M_2) hold .Then :

$$\begin{aligned}(x*y) * (z*u) &= (z*u) * (x*y) \text{ by } (M_1)\\ &= (z* (x*y))*u \text{ by } (M_2)\\ &= ((x*y)*z)*u \text{ by } (M_1)\\ &= ((x*z)*y)*u \text{ by } (M_2)\\ &= ((x*z)*u)*y \text{ by } (M_2)\\ &= (u* (x*z))*y \text{ by } (M_1)\\ &= (u*y) * (x*z) \text{ by } (M_2)\\ &= (y*u) * (x*z) \text{ by } (M_1)\\ &= (x*z) * (y*u) \text{ by } (M_1) \ .\end{aligned}$$

Therefore X is a medial of KUS-algebra . ⌂

Proposition 4.15. Let $(X; *,0)$ be a medial of KUS-algebra. Then G(X) is an KUS-ideal of X.

Proof: Since $0*0 = 0$ by (kus_3), hence $0 \in G(X)$. Next , let $x, y, z \in G(X)$ be such that $(z*y) \in G(X)$ and $(y*x) \in G(X)$, then $(z*y)*0 = (z*y)$ and $(y*x)*0 = y*x$.

$(z*x)*0 = (z*x)*[(y*x)*[(z*y)*(z*x)]]$, by $(kus_1)$and (kus_3).
$= (y*x)*[(z*x)*[(z*y)*(z*x)]]$, by (kus_4) .
$= (y*x)*(z*y)$, by proposition(4.2).
$= z*[(y*x)*y]$, by (kus_4) .
$= z*[(y*y)*x]$, by theorem $(4.14(M_2))$.
$= z*(0*x) = z*x$.

This means that G(X) is a KUS-ideal of X. This completes the proof .⌂

Corollary 4.16. Let $(X; *,0)$ be a medial of KUS-algebra. Then B(X) is a KUS-ideal of X.

Proof: Since $0*(0*0) = 0$, by corollary(4.3), $0 \in B(X)$. Let $x, y, z \in B(X)$ be such that $(z*y) \in B(X)$ and $(y*x) \in B(X)$, then $(z*y)*0 = 0$ and $(y*x)*0 = 0$.

$(z*x)*0 = (z*x)*[(y*x)*[(z*y)*(z*x)]]$, by $(kus_1)$and (kus_3).
$= (z*x)*[0*[0*(z*x)]]$
$= (z*x)*(z*x)$, by (kus_2).
$= 0$, by (kus_3) .

Therefore B(X) is a KUS-ideal of X. ⌂

Proposition 4.17. If S is a KUS-sub-algebra of a KUS-algebra $(X; *,0)$, then $G(X) \cap S = G(S)$.

Proof: It is obvious that $G(X) \cap S \subseteq G(S)$. If $x \in G(S)$, then $x*0 = x$ and $x \in S \subseteq X$. Then $x \in G(X)$ and so $x \in G(X) \cap S$, which proves the proposition. ⌂

If X is an associative KUS-algebra , then for any $x \in B(X)$,
$0 = (x*x)*x = x*(x*x) = 0*x = x$ .Thus B(X) is a zero ideal i.e., $B(X) = \{0\}$. Hence any associative KUS-algebra X is p-semi-simple.

Theorem 4.18 . The G-part $(G(X); *)$ of an associative medial of KUS-algebra X is a group in which every element is an involution .

Proof:
Let X be an associative KUS-algebra and $x, y \in G(X)$. Then $(x*y)*0 = x*y$. Hence $x*y \in G(X)$ by proposition(4.12), i.e., G(X) is closed under $(*)$. For any $x \in G(X)$, we have $x*0 = x$. By (kus_2), $0*x = x$ holds in a KUS-algebra X. Therefore $0*x = x*0 = x$ in the G-part G(X) of an associative KUS-algebra X. This means that $(G(X); *)$ is a monoid. Moreover, $x*x = 0$ shows that x has an inverse and x is an involution . Hence $(G(X); *)$ is a group which every element is an involution. ⌂

Proposition 4.19. An associative KUS-algebra X satisfying $x * 0 = x$ for any $x \in X$ is commutative, i.e., $x * y = y * x$ for any $x, y \in X$.

Proof: For any $x, y \in X$, $y * x = y * (x * 0) = x * (y * 0) = x * y$, proving the proposition .⌂

Corollary 4.20. The G-part $(G(X); *)$ of an associative medial of KUS-algebra X is an abelian group in which every element is an involution .

Proof: It follows immediately from proposition (4.19) and Theorem (4.18) . ⌂

Theorem 4.21. Let $(X; *,0)$ be a KUS-algebra. If $G(X) = X$, then X is p-semi-simple.

Proof: Assume that $G(X) = X$. Since , $\{0\} = G(X) \cap B(X) = X \cap B(X) = B(X)$. Hence X is p-semi-simple. ⌂

Theorem 4.22. If $(X; *,0)$ is a KUS-algebra of order 3, then $|G(X)| \neq 3$, that is,

$G(X) \neq X$.

Proof: For the sake of convenience, let $X = \{0,a,b\}$ be a KUS-algebra. Assume that $|G(X)| = 3$, that is, $G(X) = X$. Then $0 * 0 = 0$, $a * 0 = a$, and $b * 0 = b$. From $(kus_3)$and (kus_2), then $x * x = 0$ and $0 * x = x$, it follows that $a * a = 0$, $b * b = 0$, $0 * a = a$, and $0 * b = b$. Now let $a * b = 0$. Then 0, a, and b are candidates of the computation.
If $b * a = 0$, then $b * a = 0 = a*b$ and so $a * (b * a) = b * (a * a)$. Hence $a * 0 = b * 0$. By the cancellation law in G(X), a = b, a contradiction.
If $b * a = a$, then $a = b * a = b * (a * 0) = a * (b * 0) = a * b = 0$, a contradiction.

For the case $b * a = b$, we have $b = b * a = b * (a * 0) = a * (b * 0) = a * b = 0$, which is also a contradiction.

Next, if $a * b = a$, then $b * [(b * a) * a] = b * (a * a) = b * 0 = b \neq 0$. This leads to the conclusion that Proposition (2.7(3)) does not hold, a contradiction.

Finally, let $a * b = b$.

If $b * a = 0$, then $b = a * b = a * (b * 0) = b * (a * 0) = b * a = 0$, a contradiction.

If $b * a = a$, $b = b * 0 = b * (a * a) = a * (b * a) = a * a = 0$, a contradiction.

For the case $b * a = b$, we have $a = a * 0 = a * (b * b) = b * (a * b) = b * b = 0$, which is again a contradiction. This completes the proof. ⌂

Proposition 4.23. If (X; $*$,0) is a KUS-algebra of order 2, then in every case the G-part G(X) of X is an KUS-ideal of X.

Proof: Let |X| = 2. Then either G(X) = {0} or G(X) = X. In either case, G(X) is an ideal of X. ⌂

Theorem 4.24. Let (X; $*$,0) be a KUS-algebra of order 3. Then G(X) is an KUS-ideal of X if and only if |G(X)| = 1.

Proof: Let X := {0,a,b} be a KUS-algebra. If |G(X)| = 1, then G(X) = {0} is the trivial ideal of X.

Conversely, assume that G(X) is an ideal of X. By Theorem (4.22), we know that either |G(X)| = 1 or |G(X)| = 2. Suppose that |G(X)| = 2. Then either G(X) = {0,a}or G(X) = {0,b}. If G(X) = {0,a}, then $b * a \notin G(X)$ because G(X) is an ideal of X.

Hence $a * b = b$. Then $a = a * 0 = a * (b * b) = b * (a * b) = b * b = 0$, which is a contradiction. Similarly, G(X) = {0,b} leads to a contradiction. Therefore $|G(X)| \neq 2$ and so |G(X)| = 1. ⌂

References

[1] C. Prabpayak and U. Leerawat, On ideals and congurences in KU-algebras , scientia magna journal , 5 (2009) , no .1, 54-57 .

[2] C. Prabpayak and U. Leerawat, On isomorphisms of KU-algebras , scientia magna journal , 5 (2009) , no .3 , 25-31 .

[3]

[4] J. Meng, Ideal in BCK-algebras, Pure Appl. Math. , 2 (1986), 68-76.

[5] K. Iseki and S. Tanaka, An introduction to the theory of BCK-algebras, Math. Japon. , 23 (1978), 1-26.

[6] W. A. Dudek, On proper BCC-algebras, Bull. Inst. Math. Academia Sinica , 20 (1992), 137-150.

[7] Y. Komori, The class of BCC-algebras is not a variety, Math. Japon. , 29 (1984), 391-394.

Received: December, 2012