Robustness Inequalities for Markov Control Processes with Stochastic Discounting

E. Gordienko¹, A. Garcia² and J. Ruiz de Chavez³

Departamento de Matemáticas
Universidad Autónoma Metropolitana
Unidad Iztapalapa
Av. San Rafael Atlixco N 186
Col. Vicentina C.P. 09340, Iztapalapa
Mexico D.F.

¹gord@xanum.uam.mx

²agar@xanum.uam.mx and ³jrch@xanum.uam.mx

Abstract

We study the stability of an optimal control of general Markov chains under perturbations of the transition probabilities. The criterion of optimality is the expected total discounted cost with a random rate of discounting. We give upper bounds for the stability index which are expressed in terms of the weighted total variation distance between the transition probabilities of an original and a perturbed processes. In addition, we show how the inequalities found can be applied to estimate the robustness of the optimal control for some controlled queues and for a certain consumption-investment process.

Mathematics Subject Classification: 90C40, 90C31

Keywords: Approximating process; Discounted cost; Markov control (decision) process; Random rate; Stability index; Weighted total variation norm

1 Introduction

In this paper we investigate robustness (stability) properties of discrete time Markov control processes on Borel spaces with the expected total discounted cost as a performance index. The one-stage cost is allowed to be unbounded. The discount factors are random and can depend on their previous values and on the preceding states and actions. The interest in such control optimization

models has been arisen in the last years in some areas of applied probability such as financial mathematics, see [7, 11]. Issues of existence and approximate calculation of optimal control policies for general Markov processes with stochastic discounting were studied in the recent papers [2, 3].

We will assume that the dynamics of certain “real” control process Q , together with the evolution of the discount factor are governed by a transition probability p , which is known only approximately. Thus a decision maker, who is searching for the control policy π_* which minimizes the expected total discounted cost V for the process Q , disposes only with an approximation $\tilde{p}$ of the unknown transition probability p , which is obtained by some statistical estimations or theoretical considerations.

The transition probability $\tilde{p}$ determines the approximating control process $\tilde{Q}$ with the expected total discounted cost denoted by $\tilde{V}$. Let us suppose that the control policy $\tilde{\pi}_*$ that minimizes $\tilde{V}$ can be found. Considering $\tilde{\pi}_*$ as an available approximation to the unknown policy π_* , we evaluate the quality of such approximation by mean of the following *stability index*:

$$\Delta = V(\tilde{\pi}_*) - V(\pi_*) \geq 0.$$

The value of Δ represents the excess of the cost paid when the policy $\tilde{\pi}_*$ is applied to control the original process Q instead of the best (optimal) policy π_* .

In this paper we extend the technique used for the evaluation of the stability index for Markov control chains with a fixed discount factor, see [4]. As a result we obtain robustness inequalities of the form:

$$\Delta \leq Md(p, \tilde{p}), \quad (1.1)$$

where M is an explicit function of an initial state of a process, and d is certain “distance” between the transition probabilities. The distance d in (1.1) is defined in terms of the weighted total variation norm and, partially through the Kantorovich metric. The inequalities such as (1.1) yield to the stability of the control optimization problem in the sense that if $d(p, \tilde{p}) \rightarrow 0$, then $\Delta \rightarrow 0$. In this case the policies π_* and $\tilde{\pi}_*$ provide asymptotically the same quality of control. It is worthful to mention that there are unstable total discounted cost control optimization problems, see [5].

Since the distance $d(p, \tilde{p})$ in (1.1) involves the unknown (at least partly) transition probability p , this distance can not be calculated in practice. However, in order to apply inequalities such as (1.1) it suffices to possess some available upper bound for $d(p, \tilde{p})$. Such bounds are frequently obtainable. For example, a rather typical case is when $\tilde{p}$ is a statistical estimate of p . Then some known bounds of the rate of convergence of $\tilde{p}$ to p may be used.

In the last section of the paper we discuss the application of the inequalities found to a queue system with a controlled service rate and to the problem of dynamic consumption-investment optimization over an infinite time interval.

2 Control Model and its Approximation

Let us consider two Markov control processes Q and $\tilde{Q}$ with discrete time ($t = 0, 1, 2, \dots$) which are defined on the same *state-rate space* $Y = X \times [\alpha_*, \infty)$, where X is a given Borel space where the “states” x_t , $t = 0, 1, 2, \dots$ evolve, and $[\alpha_*, \infty)$ with a given $\alpha_* > 0$ is the interval where the random “rates” α_t , $t = 0, 1, 2, \dots$ take their values. Thus both x and α are treated as a part of the state tuple Y .

Both processes Q and $\tilde{Q}$ have a common *action set* A , which is a Borel space with given subsets $A(x, \alpha)$, $(x, \alpha) \in Y$ of admissible actions.

We also suppose that for each $(x, \alpha) \in Y$ the set $A(x, \alpha)$ is *compact*, and the set

$$\mathbb{K} := \{(x, \alpha, a); (x, \alpha) \in Y, a \in A(x, \alpha)\}$$

is a Borel subset of $Y \times A$, such that $\mathbb{K}$ contains the graph of a measurable map from Y to A .

The only difference between the control processes Q and $\tilde{Q}$ is due to their different *transition probabilities*, which are denoted, respectively, by:

$$p(B|x, \alpha, a) \text{ and } \tilde{p}(B|x, \alpha, a), \quad B \in \mathcal{B}_Y \text{ (Borel } \sigma\text{-algebra)}, (x, \alpha, a) \in \mathbb{K}.$$

In what follows $\tilde{p}$ is interpreted as an *available approximation* of the unknown transition probability p of the “real” control process Q .

The *one-stage cost* $c(x, a)$, $x \in X$, $a \in \bigcup_{\alpha \in [\alpha_*, \infty)} A(x, \alpha)$ is a given measurable function that takes values in $\mathbb{R}$.

One can think of the dynamics of the process Q as follows. If at the instant t the “state” is $x_t = x$, the rate is $\alpha_t = \alpha$, and the action $a_t = a$ is chosen, then the distribution of the pair (x_{t+1}, α_{t+1}) of the “state-rate” is determined by the transition probability:

$$p((x_{t+1}, \alpha_{t+1}) \in B \mid x, \alpha, a), \quad B \in \mathcal{B}_Y.$$

In addition, at the time t the randomly discounted cost

$$\mathbf{e}^{-(\alpha_0 + \dots + \alpha_{t-1})} c(x_t, a_t)$$

is paid.

Remark 1. For the sake of brevity we will name a “state-rate” pair (x, α) simply as a state (of a process).

The control policy $\pi = \{\pi_t, t = 0, 1, 2, \dots\}$ is defined in a standard way (see, for instance, [1]), and represents a sequence of actions (controls) at successive times. In general, π_t (randomized or not) depends on the whole “history”

of the process upon the time t , but we restrict our attention to *stationary control policies* (or briefly: *stationary policies*).

A policy $f = \{f, f, \dots\}$ is called *stationary* if for each $t \geq 0$, $\pi_t \equiv a_t = f(x_t, \alpha_t)$ with $f : Y \rightarrow A$ being a measurable function such that $f(x, \alpha) \in A(x, \alpha)$ for each $(x, \alpha) \in Y$. In words, the application of a stationary policy f results in a choice of the control $a_t = f(x_t, \alpha_t)$ when the process is in the state (x_t, α_t) . This assumes that both variables x_t and α_t are completely observable.

For each given *initial state* $(x, \alpha) \in Y$ and control policy π , the canonical probability measures on the space of trajectories:

$$\{x, \alpha, a_0; x_1, \alpha_1, a_1; \dots, x_t, \alpha_t, a_t; \dots\}$$

are defined in a standard way, (see, for example [9]). In the rest of the paper the expectations with respect to these measures are denoted by $E_{x,\alpha}^\pi$ and $\tilde{E}_{x,\alpha}^\pi$ for the processes Q and $\tilde{Q}$, respectively.

Considering the process Q and $\tilde{Q}$ with an arbitrary but fixed initial state $(x, \alpha) \in Y$ and control policy π , the *expected total discounted costs* are defined as follows:

$$V_\pi(x, \alpha) := E_{x,\alpha}^\pi \left[\sum_{t=0}^{\infty} e^{-S_t} c(x_t, a_t) \right], \quad (2.1)$$

$$\tilde{V}_\pi(x, \alpha) := \tilde{E}_{x,\alpha}^\pi \left[\sum_{t=0}^{\infty} e^{-\tilde{S}_t} c(\tilde{x}_t, \tilde{a}_t) \right], \quad (2.2)$$

where

$$\begin{aligned} S_t &= \alpha + \alpha_1 + \dots + \alpha_{t-1}, & t \geq 2; & & (S_0 := 0, S_1 := \alpha), \\ \tilde{S}_t &= \alpha + \tilde{\alpha}_1 + \dots + \tilde{\alpha}_{t-1}, & t \geq 2; & & (\tilde{S}_0 := 0, \tilde{S}_1 := \alpha). \end{aligned}$$

Note that under the hypotheses of the next section the expectations in (2.1) and (2.2) are well defined and finite.

The control policies π_* and $\tilde{\pi}_*$ are *optimal* correspondingly for the process Q and $\tilde{Q}$ if for all $(x, \alpha) \in Y$

$$\begin{aligned} V_{\pi_*}(x, \alpha) &= \inf_{\pi} V_\pi(x, \alpha) =: V_*(x, \alpha), \\ \tilde{V}_{\tilde{\pi}_*}(x, \alpha) &= \inf_{\pi} \tilde{V}_\pi(x, \alpha) =: \tilde{V}_*(x, \alpha), \end{aligned}$$

where inf is taken over all possible policies of control.

The functions $V_*, \tilde{V}_* : Y \rightarrow [-\infty, \infty)$ are called the *value functions* of Q and $\tilde{Q}$, respectively.

3 Assumptions. Stability Index.

The following conditions are needed to guarantee the existence of a minimizer in an optimality equation, which in turn, yields to the existence of stationary optimal policies of control.

Assumption 1 (*Continuity conditions*)

- (a) The map $(x, \alpha) \rightarrow A(x, \alpha)$ is upper semicontinuous with respect to the Hausdorff metric.
- (b) The cost function c is lower semicontinuous.
- (c) For every bounded continuous function $u : Y \rightarrow \mathbb{R}$ the functions

$$u'(x, \alpha, a) := \int_Y u(y, \beta) p(dy d\beta | x, \alpha, a),$$

$$u''(x, \alpha, a) := \int_Y u(y, \beta) \tilde{p}(dy d\beta | x, \alpha, a)$$

are continuous on the set $\mathbb{K}$.

To handle the cases of bounded and unbounded cost functions we will use the Banach space $(\mathcal{B}_w, \|\cdot\|_w)$ defined as follows. Let $w : X \rightarrow [1, \infty)$ be a given measurable function. The set $\mathcal{B}_w$ consists of all measurable functions $u : Y \rightarrow \mathbb{R}$ such that

$$\|u\|_w := \sup_{x \in X, \alpha \in [\alpha_*, \infty)} \frac{|u(x, \alpha)|}{w(x)} < \infty. \quad (3.1)$$

Assumption 2 (*Bounding conditions*)

- (a) For all $x \in X$ and for all $a \in \bigcup_{\alpha \in [\alpha_*, \infty)} A(x, \alpha)$ it follows that $|c(x, a)| \leq w(x)$.
- (b) There exists a number $\beta_* \in [0, 1)$ such that for each $(x, \alpha, a) \in \mathbb{K}$,

$$\int_Y w(y) p(dy d\beta | x, \alpha, a) \leq e^{\alpha_*} \beta_* w(x), \quad (3.2)$$

$$\int_Y w(y) \tilde{p}(dy d\beta | x, \alpha, a) \leq e^{\alpha_*} \beta_* w(x). \quad (3.3)$$

Applying the method given in [8], it is easy to prove that under Assumptions 1 and 2 the value functions V_* , $\tilde{V}_*$ satisfy the optimality equations. Moreover from these equations, it follows the existence of optimal stationary control policies f_* and $\tilde{f}_*$ for the processes Q and $\tilde{Q}$, respectively.

The *stability index* Δ is defined as in [4, 5, 6]:

$$\begin{aligned} \Delta(x, \alpha) &:= V_{\tilde{f}_*}(x, \alpha) - V_{f_*}(x, \alpha) \\ &\equiv V_{\tilde{f}_*}(x, \alpha) - V_*(x, \alpha) \geq 0, \quad (x, \alpha) \in Y. \end{aligned} \quad (3.4)$$

4 Robustness Inequalities

Proposition 1. *Suppose that Assumptions 1 and 2 hold. Then for any initial state $(x, \alpha) \in Y$,*

$$\Delta(x, \alpha) \leq 2e^{-\alpha_*}(1 - \beta_*)^{-2}w(x)\delta_w(p, \tilde{p}), \quad (4.1)$$

where

$$\delta_w(p, \tilde{p}) := \sup_{(x, \alpha, \alpha) \in \mathbb{K}} \frac{1}{w(x)} \int_Y w(y) |p(dy d\beta | x, \alpha, a) - \tilde{p}(dy d\beta | x, \alpha, a)|,$$

and $|\nu(\cdot)|$ stands for the *total variation* of a signed finite measure ν .

Remark 2. *Let μ_ξ and μ_η be the distributions of random vectors ξ and η taking values in a Borel space Y . It is well-known that*

$$\int_Y w(z) |\mu_\xi - \mu_\eta|(dz) = \sup_{\varphi: |\varphi| \leq w} |E\varphi(\xi) - E\varphi(\eta)|.$$

When $Y = \mathbb{R}^k$ and μ_ξ, μ_η have densities ρ_ξ, ρ_η with respect to the Lebesgue measure,

$$\int_Y w(z) |\mu_\xi - \mu_\eta|(dz) = \int_{\mathbb{R}^k} w(z) |\rho_\xi(z) - \rho_\eta(z)| dz.$$

Proof: For each stationary policy $f = f(x, \alpha)$ we define the operators:

$T_f, \tilde{T}_f := \mathcal{B}_w \rightarrow \mathcal{B}_w$ by:

$$T_f u(x, \alpha) := c(x, f(x, \alpha)) + e^{-\alpha} \int_Y u(y, \beta) p(dy d\beta | x, \alpha, f(x, \alpha)), \quad (4.2)$$

$$\tilde{T}_f u(x, \alpha) := c(x, f(x, \alpha)) + e^{-\alpha} \int_Y u(y, \beta) \tilde{p}(dy d\beta | x, \alpha, f(x, \alpha)). \quad (4.3)$$

We show that both operators maps $\mathcal{B}_w$ into $\mathcal{B}_w$, and moreover they are contractive on $\mathcal{B}_w$.

By (3.1) and Assumption 2 for any $u \in \mathcal{B}_w$, we get

$$\begin{aligned} \|T_f u\|_w &\leq \sup_{x, a} \frac{|c(x, a)|}{w(x)} + e^{-\alpha_*} \sup_{x, \alpha} \frac{1}{w(x)} \int_Y |u(y, \beta)| p(dy d\beta | x, \alpha, f(x, \alpha)) \\ &\leq 1 + e^{-\alpha_*} \sup_{x, \alpha, a} \frac{\|u\|_w}{w(x)} \int_Y w(y) p(dy d\beta | x, \alpha, a) \\ &\leq 1 + \beta_* \|u\|_w. \end{aligned}$$

In addition, using (4.2) we get that for every $u, v \in \mathcal{B}_{w_*}$

$$\begin{aligned} \|T_f u - T_f v\|_w &\leq e^{-\alpha_*} \sup_{x, \alpha} \frac{1}{w(x)} \int_Y |u(y, \beta) - v(y, \beta)| p(dy d\beta | x, \alpha, f(x, \alpha)) \\ &\leq e^{-\alpha_*} \sup_{x, \alpha, a} \frac{\|u - v\|_w}{w(x)} \int_Y w(y) p(dy d\beta | x, \alpha, a) \\ &\leq \beta_* \|u - v\|_w, \end{aligned} \quad (4.4)$$

due to Assumption 2.

It is showed similarly to (4.4) that for the operators $\tilde{T}_f$ in (4.3) we have:

$$\|\tilde{T}_f u - \tilde{T}_f v\|_w \leq \beta_* \|u - v\|_w, \quad u, v \in \mathcal{B}_w. \quad (4.5)$$

Since the policies f_* and $\tilde{f}_*$ are optimal for the process Q and $\tilde{Q}$, respectively, it follows from (3.4) and (3.1) that

$$\begin{aligned} \Delta(x, \alpha) &= V_{\tilde{f}_*}(x, \alpha) - \tilde{V}_{\tilde{f}_*}(x, \alpha) + \inf_f \tilde{V}_f(x, \alpha) - \inf_f V_f(x, \alpha) \\ &\leq 2 \sup_f |\tilde{V}_f(x, \alpha) - V_f(x, \alpha)| \\ &\leq 2w(x) \sup_f \|\tilde{V}_f - V_f\|_w. \end{aligned} \quad (4.6)$$

In the proof of Theorem 10 in [2] it was shown that for every stationary policy f ,

$$\tilde{V}_f = \tilde{T}_f \tilde{V}_f \quad \text{and} \quad V_f = T_f V_f.$$

Hence by (4.5)

$$\begin{aligned} \|\tilde{V}_f - V_f\|_w &= \|\tilde{T}_f \tilde{V}_f - T_f V_f\|_w \\ &\leq \|\tilde{T}_f \tilde{V}_f - \tilde{T}_f V_f\|_w + \|\tilde{T}_f V_f - T_f V_f\|_w \\ &\leq \beta_* \|\tilde{V}_f - V_f\|_w + \|\tilde{T}_f V_f - T_f V_f\|_w. \end{aligned} \quad (4.7)$$

Therefore, in view of (4.2) and (4.3),

$$\begin{aligned} \|\tilde{V}_f - V_f\|_w &\leq \frac{1}{1-\beta_*} \|\tilde{T}_f V_f - T_f V_f\|_w \\ &\leq \frac{e^{-\alpha_*}}{1-\beta_*} \sup_{x, \alpha} \frac{1}{w(x)} \|V_f\|_w \\ &\quad \times \left| \int_Y V_f(y, p) [p(dy d\beta | x, \alpha, f(x, \alpha)) - \tilde{p}(dy d\beta | x, \alpha, f(x, \alpha))] \right| \\ &\leq \frac{e^{-\alpha_*}}{1-\beta_*} \sup_{(x, \alpha, a) \in \mathbb{K}} \frac{1}{w(x)} \|V_f\|_w \\ &\quad \times \int_Y w(y) |p(dy d\beta | x, \alpha, a) - \tilde{p}(dy d\beta | x, \alpha, a)|. \end{aligned} \quad (4.8)$$

To obtain the desired inequality (4.1) it suffices to combine (4.6), (4.8) and to observe that by (4.4), it follows that

$$\begin{aligned}\|V_f\|_w &= \|T_f V_f\|_w \leq \|T_f V_f - T_f 0\|_w + \|T_f 0\|_w \\ &\leq \beta_* \|V_f\|_w + \|T_f 0\|_w, \quad \text{or} \\ \|V_f\|_w &\leq (1 - \beta_*)^{-1} \|T_f 0\|_w \leq (1 - \beta_*)^{-1},\end{aligned}\tag{4.9}$$

where we have used Assumption 2, (a). $\square$

Now we focus on a particular class of Markov control processes with stochastic discounting for which a “state part” $\{x_t, t \geq 0\}$ and a “rate part” $\{\alpha_t, t \geq 0\}$ of trajectories evolve “independently” (in the sense expressed in (4.10) below). We introduce the following assumptions.

Assumption 3.

There exist transition probabilities $q, \tilde{q}, r$ and $\tilde{r}$ such that for every $(x, \alpha, a) \in \mathbb{K}$,

$$\begin{aligned}p(dy d\beta | x, \alpha, a) &= q(dy | x, a) r(d\beta | \alpha, a), \\ \tilde{p}(dy d\beta | x, \alpha, a) &= \tilde{q}(dy | x, a) \tilde{r}(d\beta | \alpha, a).\end{aligned}\tag{4.10}$$

Remark 3. (a) In general, equations (4.10) do not provide the independence of $\{x_t, t \geq 0\}$ and $\{\alpha_t, t \geq 0\}$ since a control can depend on both, x and α .

(b) In the right hand side of inequality (4.11) below and in Proposition 2 later, the supremum in the first term is taken over

$$(x, a) \in \{x \in X, a \in \bigcup_{\alpha \in [\alpha_*, \infty)} A(x, \alpha)\},$$

while in the second term the supremum is taken over

$$(\alpha, a) = \{\alpha \in [\alpha_*, \infty), a \in \bigcup_{x \in X} A(x, \alpha)\}.$$

Corollary 1. Suppose that Assumptions 1, 2 and 3 hold. Then for each $(x, \alpha) \in Y$

$$\begin{aligned}\Delta(x, \alpha) &\leq 2(1 - \beta_*)^{-2} w(x) \times \\ &\left\{ e^{-\alpha_*} \sup_{x, a} \frac{1}{w(x)} \int_X w(y) |q(dy | x, a) - \tilde{q}(dy | x, a)| \right. \\ &\left. + \beta_* \sup_{\alpha, a} \int_{\alpha_*}^{\infty} |r(d\beta | \alpha, a) - \tilde{r}(d\beta | \alpha, a)| \right\}.\end{aligned}\tag{4.11}$$

Proof: We bound the integral term in the middle part of the right-hand side of inequality (4.8) as follows:

$$\begin{aligned}
d &:= \left| \int_{X \times [\alpha_*, \infty)} V_f(y, \beta) q(dy|x, f(x, \alpha)) r(d\beta|\alpha, f(x, \alpha)) \right. \\
&\quad \left. - \int_{X \times [\alpha_*, \infty)} V_f(y, \beta) \tilde{q}(dy|x, f(x, \alpha)) \tilde{r}(d\beta|\alpha, f(x, \alpha)) \right| \\
&\leq \left| \int_{X \times [\alpha_*, \infty)} V_f(y, \beta) q(dy|x, f(x, \alpha)) r(d\beta|\alpha, f(x, \alpha)) \right. \\
&\quad \left. - \int_{X \times [\alpha_*, \infty)} V_f(y, \beta) \tilde{q}(dy|x, f(x, \alpha)) r(d\beta|\alpha, f(x, \alpha)) \right| \\
&\quad + \left| \int_{X \times [\alpha_*, \infty)} V_f(y, \beta) \tilde{q}(dy|x, f(x, \alpha)) r(d\beta|\alpha, f(x, \alpha)) \right. \\
&\quad \left. - \int_{X \times [\alpha_*, \infty)} V_f(y, \beta) \tilde{q}(dy|x, f(x, \alpha)) \tilde{r}(d\beta|\alpha, f(x, \alpha)) \right| \\
&=: d_1 + d_2.
\end{aligned} \tag{4.12}$$

Integrating (3.3) over $\beta \in [\alpha_*, \infty)$ we get that

$$\int_X w(y) \tilde{q}(dy|x, a) \leq e^{\alpha_*} \beta_* w(x). \tag{4.13}$$

Now we apply Fubini's Theorem to give upper bounds for d_1 and d_2 in (4.12). This theorem is fulfilled here. Indeed, for instance for the forth integral in (4.12), denoting $a = f(x, \alpha)$ we obtain (see (4.9)):

$$\begin{aligned}
&\int_{X \times [\alpha_*, \infty)} |V_f(y, \beta)| \tilde{q}(dy|x, a) r(d\beta|\alpha, a) \\
&\leq (1 - \beta_*)^{-1} \int_{X \times [\alpha_*, \infty)} w(y) \tilde{q}(dy|x, a) r(d\beta|\alpha, a) \\
&= (1 - \beta_*)^{-1} \int_{\alpha_*}^{\infty} r(d\beta|\alpha, a) \int_X w(y) \tilde{q}(dy|x, a) < \infty.
\end{aligned}$$

The boundedness of the last integral follows from the inequality (4.13). Therefore, proceeding as in (4.8) and using (4.9) we get:

$$\begin{aligned}
d_1 &= \left| \int_{\alpha_*}^{\infty} r(d\beta|\alpha, a) \int_X V_f(y, \beta) [q(dy|x, a) - \tilde{q}(dy|x, a)] \right| \\
&\leq (1 - \beta_*)^{-1} \int_X w(y) |q(dy|x, a) - \tilde{q}(dy|x, a)|.
\end{aligned} \tag{4.14}$$

On the other hand, for $a = f(x, \alpha)$,

$$\begin{aligned} d_2 &= \left| \int_X \tilde{q}(dy|x, a) \int_{\alpha_*}^{\infty} V_f(y, \beta) [r(d\beta|\alpha, a) - \tilde{r}(d\beta|\alpha, a)] \right| \\ &\leq (1 - \beta_*)^{-1} \int_X \tilde{q}(dy|x, a) \int_{\alpha_*}^{\infty} w(y) |r(d\beta|\alpha, a) - \tilde{r}(d\beta|\alpha, a)| \quad (4.15) \\ &\leq (1 - \beta_*)^{-1} e^{\alpha_*} \beta_* w(x) \int_{\alpha_*}^{\infty} |r(d\beta|\alpha, a) - \tilde{r}(d\beta|\alpha, a)| \end{aligned}$$

due to inequality (4.13). Now we apply the bounds (4.14) and (4.15) to (4.12), and then use the resulting inequality in (4.8). Then the proof is completed as in Proposition 1. $\square$

Remark 4. *The second term of bound (4.11) is not quite satisfactory since it does not provide an estimation of stability in the problem with deterministic discount factors. Indeed, let us consider the case when $q \equiv \tilde{q}$ and in the model Q , $\alpha_t = \alpha = \text{constant}$ ($t = 0, 1, 2, \dots$) and also $\tilde{\alpha}_t^{(i)} = \tilde{\alpha}_i = \text{constant}$ ($t = 0, 1, 2, \dots$) in the approximating models $\tilde{Q}_i$, $i = 1, 2, \dots$. If $\tilde{\alpha}_i \rightarrow \alpha$ as $i \rightarrow \infty$ then it is reasonable to expect that in the left-hand side of inequality (4.11), i. e. the stability index $\Delta(x, \alpha)$, tends to zero as $i \rightarrow \infty$. However, in the situation considered,*

$$\int_{\alpha_*}^{\infty} |r(d\beta|\gamma, a) - \tilde{r}(d\beta|\gamma, a)| = 2 \quad \text{for all } \gamma, \text{ provided that } \tilde{\alpha}_i \neq \alpha.$$

We can overcome this shortcoming making use of additional restrictions on the structure of the processes Q and $\tilde{Q}$, such as the ones given in Assumption 4 below.

Assumption 4

- (a) For each $(x, \alpha) \in Y$: $A(x, \alpha) = A(x)$. In other words the action set $A(x, \alpha)$ does not depend on $\alpha \in [\alpha_*, \infty)$.
- (b) Equations (4.10) hold with transition probabilities r and $\tilde{r}$ not depending on controls a . Moreover,

$$\begin{aligned} \alpha_{t+1} &= G(\alpha_t, \eta_t), \quad t = 0, 1, 2, \dots, \\ \tilde{\alpha}_{t+1} &= G(\tilde{\alpha}_t, \tilde{\eta}_t), \quad t = 0, 1, 2, \dots, \end{aligned}$$

where $\{\eta_t\}$ and $\{\tilde{\eta}_t\}$ are sequences of i. i. d. random variables.

- (c) $|G(\alpha, z) - G(\alpha', z)| \leq |\alpha - \alpha'|$ for all z and $\alpha, \alpha' \geq \alpha_*$.

We will use the Kantorovich metric in the space of probability distributions. Let μ and λ be probability measures on $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$ with the corresponding distribution functions F_μ and F_λ . The *Kantorovich distance* is

$$\ell(\mu, \lambda) := \int_{-\infty}^{\infty} |F_\mu(s) - F_\lambda(s)| ds.$$

It is well-known (see, for example [12] that if ξ and η are random variables with the corresponding marginal distributions μ and λ , then

$$\ell(\mu, \lambda) \leq E|\xi - \eta|, \quad (4.16)$$

and that $\ell(\mu_n, \lambda) \rightarrow 0$ if and only if μ_n converges to λ weakly and

$$\int_{-\infty}^{\infty} |s| dF_{\mu_n}(s) \rightarrow \int_{-\infty}^{\infty} |s| dF_{\lambda}(s), \quad \text{as } n \rightarrow \infty.$$

Particularly, if $\xi_n = c_n$, $\eta = c$ (are constants) and $c_n \rightarrow c$, then $\ell(\xi_n, \eta) \rightarrow 0$.

Proposition 2. *Suppose that Assumptions 1, 2, and 4 hold. Then for each $(x, \alpha) \in Y$,*

$$\begin{aligned} \Delta(x, \alpha) \leq & 2(1 - \beta_*)^{-2} w(x) \left\{ e^{-\alpha_*} \sup_{x,a} \frac{1}{w(x)} \int_X w(y) |q(dy|x, a) \right. \\ & \left. - \tilde{q}(dy|x, a)| + \frac{e^{\alpha_*} \beta_*^2}{1 - \beta_*} \sup_{\alpha \in [\alpha_*, \infty)} \ell[r(\cdot|\alpha), \tilde{r}(\cdot|\alpha)] \right\}. \end{aligned} \quad (4.17)$$

Remark 5. *From Assumption 4 (b), and inequalities (4.16) and (4.17), it follows that for any selection of the joint distribution of $(\eta_1, \tilde{\eta}_1)$ we obtain:*

$$\sup_{\alpha \in [\alpha_*, \infty)} \ell[r(\cdot|\alpha), \tilde{r}(\cdot|\alpha)] \leq \sup_{\alpha \in [\alpha_*, \infty)} E|G(\alpha, \eta_1) - G(\alpha, \tilde{\eta}_1)|.$$

Proof. In view of definition (3.4) of the stability index $\Delta(x, \alpha)$ we can replace in the right-hand side of inequality (4.6) $\sup$ (over all stationary policies) by

$\max_{f \in \{f_*, \tilde{f}_*\}}$, where f_* and $\tilde{f}_*$ are optimal stationary policies for the processes Q and $\tilde{Q}$, respectively. Therefore, similarly to (4.6),

$$\Delta(x, \alpha) \leq 2w(x) \max \left\{ \|\tilde{V}_{f_*} - V_{f_*}\|_w, \|\tilde{V}_{\tilde{f}_*} - V_{\tilde{f}_*}\|_w \right\}. \quad (4.18)$$

The first term under the sign of $\max$ is evaluated as in (4.7) and (4.8), obtaining:

$$\|\tilde{V}_{f_*} - V_{f_*}\|_w \leq \frac{1}{1 - \beta_*} \|\tilde{T}_{f_*} V_* - T_{f_*} V_*\|_w, \quad (4.19)$$

because of $V_{f_*} = V_*$ (value function).

The second term is bounded as follows, by (4.4) we have:

$$\begin{aligned} \|\tilde{V}_{\tilde{f}_*} - V_{\tilde{f}_*}\|_w &= \|\tilde{T}_{\tilde{f}_*} \tilde{V}_{\tilde{f}_*} - T_{\tilde{f}_*} V_{\tilde{f}_*}\|_w \\ &\leq \|T_{\tilde{f}_*} V_{\tilde{f}_*} - T_{\tilde{f}_*} \tilde{V}_{\tilde{f}_*}\|_w + \|T_{\tilde{f}_*} \tilde{V}_{\tilde{f}_*} - \tilde{T}_{\tilde{f}_*} \tilde{V}_{\tilde{f}_*}\|_w \\ &\leq \beta_* \|\tilde{V}_{\tilde{f}_*} - V_{\tilde{f}_*}\|_w + \|T_{\tilde{f}_*} \tilde{V}_* - \tilde{T}_{\tilde{f}_*} \tilde{V}_*\|_w, \end{aligned}$$

since by the definition of $\tilde{f}_*$, $\tilde{V}_{\tilde{f}_*} = \tilde{V}_*$. Consequently

$$\|\tilde{V}_{\tilde{f}_*} - V_{\tilde{f}_*}\|_w \leq \frac{1}{1 - \beta_*} \|\tilde{T}_{\tilde{f}_*} \tilde{V}_* - T_{\tilde{f}_*} \tilde{V}_*\|_w. \quad (4.20)$$

The rest of the proof is carried out along the lines of Corollary 1. However now we evaluate the first integral in (4.15) with $f = f_*$ and $f = \tilde{f}_*$ using a distinct approach.

For $f = f_*$ we get in (4.15):

$$d_2 \leq \int_X \tilde{q}(dy|x, a) \left| \int_{\alpha_*}^{\infty} V_*(y, \beta) [r(d\beta|\alpha) - \tilde{r}(d\beta|\alpha)] \right|, \quad (4.21)$$

Similarly, using (4.20) instead of (4.19) we can see that for $f = \tilde{f}_*$ in (4.15)

$$d_2 \leq \int_X \tilde{q}(dy|x, a) \left| \int_{\alpha_*}^{\infty} \tilde{V}_*(y, \beta) [r(d\beta|\alpha) - \tilde{r}(d\beta|\alpha)] \right|. \quad (4.22)$$

Now we show that for every $x \in X$ the functions $V_*(x, \cdot)$, $\tilde{V}_*(x, \cdot)$ satisfy the Lipschitz condition on $[\alpha_*, \infty)$ with the constant

$$L = \frac{\beta_*}{(1 - \beta_*)^2} w(x). \quad (4.23)$$

Let us realize it, for example for V_* . In [2], it was shown that

$$V_* = TV_*, \quad (4.24)$$

where for $u \in \mathcal{B}_w$, $(x, \alpha) \in Y$,

$$Tu(x, \alpha) := \min_{a \in A(x)} \left\{ c(x, a) + e^{-\alpha} \int_{X \times [\alpha_*, \infty)} u(y, \beta) q(dy|x, a) r(d\beta|\alpha) \right\}. \quad (4.25)$$

Similarly to (4.4), it is shown that T is a contractive operator on $(\mathcal{B}_w, \|\cdot\|_w)$ with modulus β_* . Therefore in view of (4.24),

$$V_* = \lim_{n \rightarrow \infty} V_n, \quad \text{where} \quad V_n := T^n 0, \quad n \geq 0. \quad (4.26)$$

We verify by induction that for every $n \geq 0$, $x \in X$; $\alpha, \alpha' \geq \alpha_*$,

$$|V_n(x, \alpha) - V_n(x, \alpha')| \leq \frac{\beta_*}{1 - \beta_*} w(x) L_n |\alpha - \alpha'|, \quad (4.27)$$

where

$$L_n := \sum_{k=0}^{n-1} \beta_*^k \leq (1 - \beta_*)^{-1}. \quad (4.28)$$

By (4.25) and Fubini's theorem we get:

$$\begin{aligned} V_{n+1}(x, \alpha) &= \min_{a \in A(x)} \left\{ c(x, a) + e^{-\alpha} \int_X q(dy|x, a) \int_{\alpha_*}^{\infty} V_n(y, \beta) r(d\beta|\alpha) \right\} \\ &= \min_{a \in A(x)} \left\{ c(x, a) + e^{-\alpha} \int_X q(dy|x, a) EV_n[y, G(\alpha, \eta)] \right\}. \end{aligned} \quad (4.29)$$

Where η is a random variable distributed as η_t in Assumption 4, (b). In (4.29) E denotes the expectation with respect to η .

The use of Fubini's theorem in the above integrals is justified in the same way as in the proof of Corollary 1, provided that we bound $\|V_n\|_w$ as follows. In view of (4.25) and Assumption 2, repeating the calculations in the proof of Proposition 1, we obtain:

$$\|V_n\|_w \leq 1 + \beta^* \|V_{n-1}\|_w, \quad n \geq 1.$$

Iterating these inequalities we see that:

$$\|V_n\|_w \leq (1 - \beta_*)^{-1}, \quad n \geq 1. \quad (4.30)$$

Now, assuming (4.27) and using (4.29), the version of (4.13) for q , Assumption 4 (c) and (4.30), we can write for $\alpha, \alpha' \geq \alpha_*$:

$$\begin{aligned} &|V_{n+1}(x, \alpha) - V_{n+1}(x, \alpha')| \\ &\leq \max_{a \in A(x)} |e^{-\alpha} - e^{-\alpha'}| \left| \int_X q(dy|x, a) EV_n[y, G(\alpha, \eta)] \right| + \\ &\max_{a \in A(x)} \left| e^{-\alpha'} \int_X q(dy|x, a) \{EV_n[y, G(\alpha, \eta)] - EV_n[y, G(\alpha', \eta)]\} \right| \\ &\leq \max_{a \in A(x)} |\alpha - \alpha'| e^{-\alpha_*} \|V_n\|_w \int_X w(y) q(dy|x, a) \\ &+ e^{-\alpha_*} \max_{a \in A(x)} \int_X q(dy|x, a) \frac{\beta_*}{1 - \beta_*} w(y) L_n |\alpha - \alpha'| \\ &\leq \frac{\beta_*}{1 - \beta_*} w(x) [1 + \beta_* L_n] |\alpha - \alpha'|. \end{aligned} \quad (4.31)$$

But by (4.28), $1 + \beta_* L_n = L_{n+1}$, and therefore from (4.31) the inequality (4.27) follows for V_{n+1} .

In view of (4.26) V_n converges to V_* in $\mathcal{B}_w$ and therefore pointwise. Hence the function $V_*(x, \cdot)$ satisfies the Lipschitz condition with the constant L given in (4.23). The same property of $\tilde{V}_*(x, \cdot)$ from (4.22) is proved analogously.

Returning to (4.21) we can write:

$$d_2 \leq \sup_{x, a} \int_X w(y) q(dy|x, a) \left| \int_{\alpha_*}^{\infty} \frac{V_*(y, \beta)}{w(y)} [r(d\beta|\alpha) - \tilde{r}(d\beta|\alpha)] \right|, \quad (4.32)$$

where for each $y \in X$ the function $\varphi_y(\cdot) := \frac{V_*(y, \cdot)}{w(y)}$ satisfies on $[\alpha_*, \infty)$ the Lipschitz condition with the constant $\beta_*(1 - \beta_*)^{-2}$.

On the other hand, it is known (see [12]) that for distributions μ, λ , the Kantorovich distance can be expressed as follows:

$$\ell(\mu, \lambda) = \sup_{\varphi \in Lip} \left| \int_{\mathbb{R}} \varphi d\mu - \int_{\mathbb{R}} \varphi d\lambda \right|,$$

where $Lip := \{\varphi : \mathbb{R} \rightarrow \mathbb{R} : |\varphi(s) - \varphi(t)| \leq |s - t|, s, t \in \mathbb{R}\}$.

Consequently, from (4.32) it follows that:

$$\begin{aligned} d_2 &\leq \frac{\beta_*}{(1-\beta_*)^2} \ell[r(\cdot|\alpha), \tilde{r}(\cdot|\alpha)] \sup_{x,a} \int_X w(y) \tilde{q}(dy|x, a) \\ &\leq \frac{e^{\alpha_*} \beta_*^2}{(1-\beta_*)^2} w(x) \ell[r(\cdot|\alpha), \tilde{r}(\cdot|\alpha)]. \end{aligned} \quad (4.33)$$

Applying the same arguments one can see that the right-hand side of inequality (4.22) has the same upper bound as in (4.33).

To complete the proof of inequality (4.17) it suffices to carry out the following steps, which basically have been done in the proof of Corollary 1:

- (1) Start with inequalities (4.18)-(4.20).
- (2) Evaluate, say $\|\tilde{T}_{f_*} V_* - T_{f_*} V_*\|_w$, proceeding as in (4.12) and move to the definition of d_2 in (4.15), with V_* and $\tilde{V}_*$ instead of V_f .
- (3) Bound d_2 in (4.21) and (4.22). Here one can use inequality (4.33).
- (4) Finish the proof as in Corollary 1. □

5 Applications

5.1 $GI|D|1|_\infty$ queueing system with controllable service rate.

Let us consider the following control process:

$$x_{t+1} = \max\{0, x_t + a_t - \xi_t\}, \quad t = 0, 1, 2, \dots, \quad (5.1)$$

where $x_t \in X = [0, \infty)$ is the waiting time of the t -th customer ($t = 0, 1, 2, \dots$), whereas ξ_t denotes the interarrival time between the t -th and the $(t+1)$ -th customers. The service time of the t -customer a_t is nonrandom, but it is controllable in the sense that for each customer a_t is chosen from a given set $A(x_t, \alpha_t)$. We suppose that $A(x, \alpha) \equiv A$ with A being any given compact

subset of a finite interval $(0, \theta]$ with $\theta \in A$; and that $\xi_1, \xi_2, \dots$ are nonnegative i.i.d. random variables.

We assume that the dynamics of random rates is given by the following autoregressive process:

$$\alpha_{t+1} = \gamma\alpha_t + \eta_t, \quad t = 0, 1, 2, \dots, \quad (5.2)$$

where $\gamma \in [0, 1]$ is a constant, $\alpha_0 = \alpha \geq \alpha_*$ for some fixed $\alpha_* > 0$, and $\eta_1, \eta_2, \dots$ are i.i.d. random variables with values in the interval $[\alpha_*(1 - \gamma), \infty)$. Then $\alpha_t \geq \alpha_*$ for all $t \geq 0$. We assume that $\{\eta_t, t \geq 0\}$ is independent of $\{\xi_t, t \geq 0\}$.

Along with the “real” control process given in (5.1) and (5.2) (supposingly unknown distributions of $\xi := \xi_1$ and of $\eta := \eta_1$) we also consider its approximation:

$$\tilde{x}_{t+1} = \max\{0, \tilde{x}_t + a_t - \tilde{\xi}_t\} \quad t = 0, 1, 2, \dots, \quad (5.3)$$

$$\tilde{\alpha}_{t+1} = \gamma\tilde{\alpha}_t + \tilde{\eta}_t, \quad t = 0, 1, 2, \dots, \quad (5.4)$$

which, in general differ from (5.1), (5.2) in the distributions of the random variables $\tilde{\xi} := \tilde{\xi}_1, \tilde{\eta} := \tilde{\eta}_1$. They are supposed to be known by a controller. We denote by F_η and $F_{\tilde{\eta}}$ the distribution functions of η and $\tilde{\eta}$ respectively.

Assumption 5.

- (a) $E\xi > \theta$ and $E\tilde{\xi} > \theta$, (ergodicity condition).
- (b) The random variables ξ and $\tilde{\xi}$ have bounded continuous densities on $(0, \infty)$ denoted by ρ_ξ and $\rho_{\tilde{\xi}}$, respectively.

In [4], it was proved that there exists a number $d > 0$ such that if $w(x) = b(e^{dx} + \gamma), x \in [0, \infty)$, where b is arbitrary, but γ is specially chosen, then conditions (3.2) and (3.3) hold true, provided an appropriate selection of $\beta_* < 1$. It is also shown there how to calculate d .

Assumption 6. The one-stage cost c is a continuous nonnegative function such that for some $b \geq 1$,

$$c(x, a) \leq be^{dx}, \quad x \in [0, \infty), \quad a \in A.$$

Proposition 3. *If the Assumptions 5 and 6 hold, then for every $x \in [0, \infty)$, $\alpha \geq \alpha_*$,*

$$\begin{aligned} \Delta(x, \alpha) \leq & \frac{2b}{(1-\beta_*)^2}(e^{dx} + \gamma) \left\{ e^{d\theta} \int_0^\infty |\rho_\xi(s) - \rho_{\tilde{\xi}}(s)| ds \right. \\ & \left. + \frac{e^{\alpha_*} \beta_*^2}{1-\beta_*} \int_{\alpha_*(1-\gamma)}^\infty |F_\eta(s) - F_{\tilde{\eta}}(s)| ds \right\}, \end{aligned} \quad (5.5)$$

where β_* is an arbitrary, but fixed number from the interval $(e^{-\alpha_*}, 1)$ and $\gamma = (\beta_* e^{\alpha_*} - 1)^{-1}$.

To prove (5.5) it is enough to observe the following:

- (1) Assumption 1 is satisfied because of Assumption 6 and Assumption 5, (b).
- (2) As we have mentioned, Assumption 2 is also assured.
- (3) The verification of Assumption 4 is straightforward.
- (4) Hence inequality (4.17) from Proposition 2 can be applied.
- (5) In [4] it is proven that for $w(x) = b(e^{dx} + \gamma)$, the first summand on the right-hand side of (4.17) can be bounded by the first summand on the right-hand side in (5.5).
- (6) Concerning the second term in (4.17), by (5.2), (5.4) we see that for every $\alpha \geq \alpha_*$,

$$\ell(\gamma\alpha + \eta, \gamma\alpha + \tilde{\eta}) = \int_{\alpha_*(1-\gamma)}^{\infty} |F_{\eta}(s) - F_{\tilde{\eta}}(s)| ds.$$

5.2 Discrete-time model of portfolio-consumption optimization

The stability of such model with a fixed discount factor was studied in [6]. Here we outline a possible application of the stability inequality (4.17) in the extended version of the model with a random rate of discounting (for example, given by equations (5.2), (5.4)).

In this model, “the original wealth process” is defined by the equations:

$$x_t = (1 - a_0)x_{t-1} \sum_{k=1}^m a_k \zeta_t^{(k)}, \quad t = 1, 2, \dots, \quad (5.6)$$

while its perturbed version is

$$\tilde{x}_t = (1 - a_0)\tilde{x}_{t-1} \sum_{k=1}^m a_k \tilde{\zeta}_t^{(k)}, \quad t = 1, 2, \dots \quad (5.7)$$

Here the sequences of independent random vectors $\xi_t := (\zeta_t^{(1)}, \dots, \zeta_t^{(m)})$, $t \geq 1$ and $\tilde{\xi}_t := (\tilde{\zeta}_t^{(1)}, \dots, \tilde{\zeta}_t^{(m)})$ represent return rates of m different assets (“risky” in general, but the k -asset is “risky free” if the distribution of $\zeta_t^{(k)}$ is degenerated). The vector $(a_0, a_1, \dots, a_m)$ of actions (controls) belonging to certain compact subset of $\mathbb{R}^{m+1}$ denotes an allocation of the current capital x_{t-1} among a consumption $a_0 x_{t-1}$ and the investment in m asserts (“portfolio”).

The criterion of optimality of the consumption-investment policy $\pi = \{\bar{a}_t = (a_0^{(t)}, a_1^{(t)}, a_2^{(t)}, \dots, a_m^{(t)}), t \geq 1\}$ is the expected total randomly discounted cost defined in (2.1), and in (2.2) for the approximating process. (See, e.g. [10] for continuous time control problems of this kind with a fixed discount factor.) The one stage costs is defined as some sort of “generalized logarithmic utility”:

$$c(x, a) = -g[\ln(a_0 x); a_1, \dots, a_m]$$

where $g : \mathbb{R}^{m+1} \rightarrow \mathbb{R}$ is a certain continuous function.

The model turns out to be more tractable by taking the logarithm in (5.6) and (5.7). This provides a linear control processes. In [6], it was shown how, under certain mild restrictions, to construct a “Lyapunov function” $w(x)$, $x \geq 0$ to meet Assumptions 1 and 2 for these processes.

On the other hand, making suitable assumptions about the rate processes $\{\alpha_t, t \geq 0\}$, $\{\tilde{\alpha}_t, t \geq 0\}$, (for example such as in (5.2), (5.4)) we can guarantee the fulfilment of Assumption 4. Consequently, one can use the robustness inequality (4.17) to bound the stability index for the consumption-investment optimization problem discussed above. We prefer to leave out the calculations and remark only that under hypotheses (5.2), (5.4) “the rate term” in the stability inequality would be the same as in (5.5), while assuming the absolute continuity of ξ_t and $\tilde{\xi}_t$, it is possible to obtain something resembling the first term in (5.5). Unfortunately if for some k , $\zeta_t^{(k)}$ is constant (a “non risky” investment), we must admit that $\tilde{\zeta}_t^{(k)} = \zeta_t^{(k)}$ to get a meaningful application of inequality (4.17).

References

- [1] E.B. Dynkin, A.A. Yushkevich, *Controlled Markov processes*, Springer, New York, 1979.
- [2] J. González-Hernández, R.R. López-Martínez, J.R. Pérez-Hernández, Markov control processes with randomized discounted cost, *Math. Methods Oper. Res.*, **65** (2007), 27 - 44.
- [3] J. González-Hernández, R.R. López-Martínez, J.A. Minjarez-Sosa, Approximation, estimation and control of stochastic systems under a randomized discounted cost criterion, *Kybernetika*, **45** (2009), 737 - 754.
- [4] E.I. Gordienko, F.S. Salem, Robustness inequality for Markov control processes with unbounded costs, *Systems Control Lett.*, **33** (1998), 125 - 130.
- [5] E.I. Gordienko, F.S. Salem, Estimates of stability of Markov control processes with unbounded costs *Kybernetika*, **36** (2000), 195 - 210.

- [6] E.I. Gordienko, E. Lemus-Rodríguez and R. Montes-de-Oca, Discounted cost optimality problem: stability with respect to weak metrics, *Math. Methods Oper. Res.*, **68** (2008), 77 - 96.
- [7] S. Haberman, J.H. Sung, Optimal pension funding dynamics over infinite control horizon when stochastic rates of return are stationary, *Insurance Math. and Econom.*, **36** (2005), 103 - 116.
- [8] O. Hernández-Lerma and J.B. Lasserre, *Further topics on discrete-time Markov control processes*, Springer-Verlag, New York 1999.
- [9] K. Hinderer, Foundations of non-stationary dynamical programming with discrete time parameter, *Lecture Notes in Operation Research*, **33** Springer, Berlin 1970.
- [10] R. Korn, E. Korn, *Option pricing and portfolio optimization*. In: Modern methods of financial mathematics. American Mathematical Society, Providence, 2001.
- [11] R.G. Newell, W.A. Pezer, Discounting the distant future: how much do uncertain rates increase valuation?, *J. Environmental Economics and Management*, **46** (2003) 52 - 71.
- [12] S.T. Rachev and L. Rüschendorf, *Mass Transportation Problem, vol II*, Applications. Springer, New York 1998.

Received: September, 2010