

Influence of Sodium Chloride on Physiological, Biochemical and Yield Indicators of Wheat and Maize Genotypes

M. A. Khanishova, K. R. Tagieva and I. V. Azizov

Institute of Molecular Biology and Biotechnologies
Izzat Nabiyeu 11, AZ1073, Baku, Azerbaijan

This article is distributed under the Creative Commons by-nc-nd Attribution License.
Copyright © 2023 Hikari Ltd.

Abstract

Study the effect of sodium chloride on the physiological and biochemical characteristics of wheat and maize genotypes, to identify varieties tolerant to salinity. To create new salt-tolerant varieties of wheat and maize, a comparative analysis of the physiological and biochemical parameters of the parental and hybrid plants was carried out. Physiological parameters such as chlorophyll content and relative water content have been studied in parental forms and hybrids. When studying salt-tolerance of hybrids and parental forms, differences were detected in the relative amounts of chlorophyll (a+b), carotenoids and RWC.. The effect of salt on the amount of chlorophyll a, chlorophyll b, and carotenoids, which are the main physiological indicators, is manifested in different ways in both hybrids and parental forms. Based on the physiological indices determined during the grain filling stage, the highest tolerance was manifested by the parental forms, Garabagh, Gobustan, and the hybrids, ♀Garabagh×♂Gobustan, ♀Garabagh×♂Mirbashir, ♀Garabagh×♂Sharg. Under the action of salt in all genotypes of corn found a slight increase in the content of soluble sugars. The highest sugar content was observed in the hybrid Zagatala 68 x Gurur.

Keywords: wheat, maize, salinity, hybrid, genotype, chlorophyll, carotenoid, tolerance

Introduction

Salinity is one of the abiotic stress factors decreasing plant productivity. The

salinization of soils over time is particularly dangerous. The limitation of agricultural and fertile lands is an obstacle to meeting the food requirements of the population [Khan, et al, 2010]. In particular, the rapid growth of the population and the need in ensuring food security make more urgent the development of salt-tolerant varieties capable to grow in saline soils, and their extensive use. According to rough estimates, 521,700 hectares of plains in the Azerbaijan Republic were in a saline state in [Azizov, 2002]. In 2007, this parameter increased to 661.9 thousand hectares and accounted for 46.6 % of the land [Mammadov, 2007].

One of the most effective measures taken to achieve high productivity under stress is the development of plants capable to adapt to salinity. The expression of genes regulating stress tolerance increases under high salt concentrations and ensures salt tolerance of plants [Garraat, et al, 2002].

According to some authors, developing more plastic wheat varieties, suitable for the regions of the republic is required because of the disturbance of ecological balance and the presence of abiotic stress factors. Therefore, stress tolerance in plant breeding is of great importance [Rustamov, et al, 2017]. Currently, in our country, extensive research has been carried out on salt-tolerance of local wheat varieties as well as brought from abroad Huseynova, et al, 2008]. Thus, numerous studies conducted in the world and in our country showed the perspectives of the development of the wheat varieties adapted to salinity.

The purpose of the research was to study the effect of sodium chloride on the physiological and biochemical characteristics of wheat and corn genotypes, to identify varieties tolerant to salinity.

Materials and methods

The objects of the study were parental and hybrid wheat forms grown under normal and salinity (0,98 % NaCl) conditions. The objects of the study were the genotypes of wheat Gobustan, Mirbashir-128, Barakatli-95, Garabagh, Gyrmzygul, Sharg and their generation hybrids Garabagh×Gobustan, Gobustan×Barakatli-95, Gobustan×Gyrmzygul, Barakatli-95×Gobustan, Gobustan×Garabagh, Garabagh×Mirbashir, Garabagh×Sharg.

The second generation (F₂) hybrids obtained from seven combinations during the study were planted on the field of 1 m², under normal conditions, on October 25, 2019. For the experiments, 300 g NaCl was added to the soil before sowing during tillering, earing, and grain-filling stages. Sowing was conducted following the scheme P♀-F₂-P♂ 5×25 cm. Watering was performed at the tillering, earing, and grain-filling stages. Agrotechnical care was conducted in the experimental field. Samples were taken at the grain filling stage of the vegetation. Measurements were done *in vivo* and *in vitro* for the comparative study of the changes in physiological parameters caused by the salt effect.

The object of research was also the genotypes of maize Zagatala-420, Zagatala-514, Zagatala-68, Gurur and first generation F₁ hybrid Zagatal-68×Gurur. Plant seeds germinated under laboratory conditions in Petri dishes and pots with soil using 150 mmol of sodium chloride solution. Germination energy was

determined by counting three days old seedlings, and seed germination ability by counting seven days old seedlings as a percentage. In two-week old seedlings determined the content of soluble sugars.

0.1 g of leaf samples of hybrid and parental forms taken from plants grown under both normal and saline conditions were homogenized using a pestle and mortar in 96 % alcohol by adding CaCO_3 , centrifuged at 200 g, and a pure extract of chlorophyll pigments was obtained. The optical density of a solution of chlorophyll in alcohol was measured on an SP-2000 spectrophotometer at 665, 649, 440 nm, and the amounts of chlorophyll and carotenoids were determined [Wintermans, De Mots, 1965]. Water loss in leaves relative water content (RWC) was determined based on the method of Tambussi and colleagues [Tambussi, et al, 2005]. Determination of soluble sugars carried out by the anthrone-sulfuric acid method [Azizov, et al, 2019].

Data analysis and statistical analysis were conducted using Microsoft Excel. Statistical analysis was performed with the aid of the Statgraphics Plus 5.1 statistical package. The means of values were compared by Duncan's multiple range test ($p=0.05$).

Results and discussion

To develop new, pure lines, salt tolerance of hybrids and parental forms was assessed based on physiological indices in the grain filling stage of the vegetation. When studying salt-tolerance of hybrids and parental forms, differences were detected in the relative amounts of chlorophyll (a+b), carotenoids, RWC, as well as the photochemical activity of chloroplasts. The effect of salt on the amount of chlorophyll a, chlorophyll b, and carotenoids, which are the main physiological indicators, is manifested in different ways in both hybrids and parental forms. Based on the changes in the amount of pigments under salinity, the parental forms Garabagh, Gobustan, and Sharg were more tolerant. Thus, at least a 10 %-decrease in the amount of chlorophyll (a+b) pigments due to the effect of salt was observed in these parental genotypes. Whereas, the most growth in carotenoids amounted to 4–7 % in these varieties. This process can be attributed to the increase in the protective function of carotenoids under salt stress. Based on RWC values, Garabagh and Gobustan varieties had the highest water content. Water loss in these varieties did not decrease but increased by 3 %. The results obtained are presented in **Table 1**.

Based on the amounts of chlorophyll (a+b), the hybrids, ♀Garabagh×♂Gobustan, ♀Garabagh×♂Mirbashir, ♀Garabagh×♂Sharg are more tolerant to salinity. An increase in the amount of carotenoids was also observed in these hybrids. The highest values for RWC were found in the hybrid forms, ♀Garabagh×♂Gobustan, ♀Barakatli-95×♂Gobustan, ♀Garabagh×♂Sharg. RWC was 10 % higher in these hybrids compared to others under stress. A 20 %-decrease in RWC was observed in the hybrid form, ♀Gobustan×♂Gyrmyzygul.

Table 1. Physiological indices (photosynthetic pigments, RWC) of the parental and hybrid forms of wheat under 1 % NaCl* (mg/g fresh weight)

Wheat varieties	Chl(a+b)		Carotenoids		RWC	
	Control	NaCl	Control	NaCl	Control	NaCl
Gobustan	3.47	2.42	1.04	0.54	70.9	73.1
Mirbashir-128	2.95	2.03	1.03	0.95	75.5	74.9
Barakatli-95	3.29	1.61	1.08	0.97	71.9	74.2
Garabagh	2.55	2.71	1.05	0.69	89.5	83.6
Gyrmyzygul	3.57	1.56	0.87	0.74	88.9	80.8
Shark	3.38	1.50	1.17	0.38	70.5	76.9
Garabagh×Gobustan	2.98	1.50	1.17	0.51	67.9	84.1
Gobustan×Barakatli-95	2.93	2.49	1.04	0.90	98.6	72.8
Gobustan×Gyrmyzygul	2.54	2.24	0.86	1.06	75.5	79.6
Barakatli-95×Gobustan	3.33	1.51	1.35	0.62	67.4	81.8
Gobustan×Qarabagh	2.09	1.63	0.60	0.63	92.3	75.6
Garabagh×Mirbaşir-128	2.26	1.75	0.57	0.29	94.8	79.5
Garabagh×Shark	3.55	1.42	1.72	0.33	74.4	83.5

Note: * Each value represents the mean \pm SD (standard deviation) for the mean $n=3$ independent experiments $p=0.05$

The first indicator of fluorescence in varieties grown in a saline environment is light energy. Energy loss occurs under salinity. In this case, the activity of the photosystem is reduced, which leads to the weakening development of the plant.

Based on the PS II activity, the parental forms, Garabagh, Gobustan, and the hybrids, ♀Gobustan×Barakatli-95, ♀Gobustan×♂Garabagh, and ♀Gobustan×♂Sharg are more salt-tolerant.

One of the most useful indicators of wheat is grain yield. High salt concentrations contributed to a decrease in grain yield. Among the genotypes, according to this indicator, the most salt tolerant were the hybrids Barakatli-95×Gobustan and Garabakh×Mirbashir-128 (**Table 2**).

It revealed that under the influence of salt, germination and germination energy of maize seeds reduced. With the increase of sodium chloride concentration decreased the photochemical activity of chloroplasts and the content of photosynthetic pigments in seedlings.

Salinity could affect chlorophyll concentration of leaves through inhibition of synthesis of chlorophyll or an acceleration of its degradation. Impairment of the carboxylation capacity, which in turn inhibits electron transport, is indicated by the measurements of chlorophyll fluorescence. A reduced quantum yield may result from a structural impact on PS II although some authors [Lu,et al,2002] found PS II to be highly resistant to salinity stress. Salinity has been concluded to affect reaction centers of PS II either directly or via an accelerated senescence. High external salt concentrations could affect thylakoid membranes by disrupting lipid bilayer or lipid-protein associations and thus, impair electron transport activity. The efficiency of the photochemical conversion of the PS II energy decreased with increasing salt concentrations. Some authors indicate the decrease of the root

system function in plants exposed to salt stress. They assumed a more important role of toxic effects of ions [Wang, et al, 2012].

Table 2. Effect of NaCl on elements of the yield structure of durum and bread winter wheat varieties under the conditions of a growing experiment

wheat varieties	Plant height, sm		number of grains per ear		grain weight, g		crop losses, %
	control	NaCl	control	NaCl	control	NaCl	
Gobustan	90±2	68±1	62±3	45±4	3,6	1,4	62
Mirbashir-128	105±4	73±1	42±2	18±1	4,6	1,2	74
Barakatli-95	85±2	75±2	68±3	29±2	3,5	1,7	52
Garabagh	80±2	62±1	41±1	32±2	2,2	1,5	32
Gyrmyzygul	72±1	57±2	46±2	22±1	3,1	2,4	23
Shark	110±4	70±3	32±2	19±1	4,2	1,4	67
Garabagh×Gobustan	120±3	65±2	52±3	36±2	1,4	1,1	22
Gobustan×Barakatli-95	76±2	100±4	32±2	20±1	2,8	2,2	22
Gobustan×Gyrmyzygul	118±4	70±2	34±1	27±2	2,9	2,3	21
Barakatli-95×Gobustan	78±1	63±1	40±2	21±1	1,9	1,6	10
Gobustan×Qarabagh	80±1	60±2	38±1	34±2	4,2	2,8	34
Garabagh×Mirbaşir-128	90±2	60±1	54±3	33±3	1,9	1,7	11
Garabagh×Shark	90±2	70±2	60±2	22±2	3,2	1,3	60

Note: * Each value represents the mean ±SD (standard deviation) for the mean n=3 independent experiments p=0.05.

Under the conditions of salt action, a slight increase in the content of soluble sugars was found in all genotypes of maize (Fig.).




Fig. Influence of sodium chloride on content sugars in seedlings of maize varieties

In general, the growth and development of plants depend on the process of photosynthesis in their green organs. Therefore, environmental stressors affecting photosynthesis also affect growth and development [Villora, et al, 1997]. A positive correlation between the rate of photosynthesis and productivity has been found in various plants under salinity [Perez, et al, 1996].

The decrease in RWC due to stress indicates that the cell does not have the turgor necessary for the tension process to take place [Katterji, et al, 1997]. The response of plants to stressors is different, depending on the genetic material. Thus, the genetic material regulates the speed and consistency of protein synthesis required under stress. Some difficulties in the cultivation of salt-tolerant forms are attributed to the complexity and polygenic nature of genes. It is known that under the salt stress, the external water potential decreases, the absorption of biogenic metal ions by the roots becomes difficult, and the chlorine and sodium ions have a toxic effect on plant metabolism. These three possible effects of salt stress have a detrimental effect on plant growth, development and yield. [Muhammad, et al, 2015; Munns, et al, 2006]. Osmotic stress is associated with the accumulation of ions in the soil solution, while malnutrition and the specific effects of ions are associated to the accumulation of ions, mainly sodium and chloride, to toxic levels which inhibits the availability of other important elements such as calcium and potassium. Toxic levels of sodium in plant organs damage biological membranes and subcellular organelles, reducing growth and causing abnormal development before plant death. Several physiological processes, such as photosynthesis, respiration, starch metabolism and fixation of nitrogen also disrupted in salt conditions, which leads to a decrease in crop productivity. In response to this, the plant synthesizes low molecular weight solutes, including soluble carbohydrates for better absorption of water during salinity. Genotypes with a powerful genetic apparatus cope with this task and grow well in salt conditions. In the process of evolution, protective mechanisms against environmental stressors are formed in all organisms, including plants. Therefore, when assessing tolerance to stress factors, it is necessary to consider the individual characteristics of each plant genotype [9]. NaCl (0.98 %) in the soil has been found to affect physiological parameters of parental and hybrid forms differently.

Conclusion

Negative effects of salt stress were observed in the content of photosynthetic pigments, photochemical activity of PS II and RWC in parental and hybrid forms. The content of chl a, chl b, the activity of PSII, RWC and grain yield were higher in the parental forms, Garabagh, Gobustan, Barakatli-95, and in the hybrids, ♀Garabagh×♂Gobustan, ♀Barakatli-95×♂Gobustan, ♀Gobustan×♂Gyrmyzygul-1 and ♀Garabagh×♂Mirbashir-128. Due to these advantages, using these varieties for future research on the development of salt-tolerant forms can be considered expedient.

Under the conditions of salt action, a slight increase in the content of soluble sugars was found in all genotypes of maize. The highest sugar content was observed

in the hybrid Zagatala 68 x Gurur.

References

- [1] Azizov, G.Z., *Classification of the Azerbaijan Saline Soils Based on the Salinity Degree*, Baku, Elm, 2002, 29.
- [2] Azizov, I.V., Khanishova, M.A., Tagiyeva, K.R., and Gasimova, F.I., Comparative study of physiological and biochemical characteristics of hybrids and parental forms of wheat under drought. The role of physiology and biochemistry in the introduction and selection of agricultural plants, *Collection Of Materials Of The V International Scientific-Methodical Conference. Moscow*, **2** (2019), 26-29.
- [3] Garratt, L.C., Janagoundar, B.S., Lowe, K.C., Anthony, P., Power, J.B., and Davey, M.R., Salinity tolerance and antioxidant status in cotton cultures, *Free Radicle Biology and Medicine*, **33** (2002), 502-511.
[https://doi.org/10.1016/s0891-5849\(02\)00838-9](https://doi.org/10.1016/s0891-5849(02)00838-9)
- [4] Huseynova, I.M., Suleymanov, S.Yu., Azizov, I.V., Rustamova, S.M., Magerramova, E.G. and Aliev, J.A. *Effects Of High Concentrations Of Sodium Chloride On Photosynthetic Membranes Of Wheat Genotypes*, Scientific works of the Institute of Botany of ANAS., 2008, vol. XXYIII, 230-238. (in Russian)
- [5] Katerji, N., Van Hoorn, J. W., Hamdy, A., Mastrorilli, M., and Mou Karzel, E. Osmotic adjustment of sugar beets in response to soil salinity and its influence on stomatal conductance, growth, and yield, *Agricul. Water Manage.*, **34** (1997), 57-69. [https://doi.org/10.1016/s0378-3774\(96\)01294-2](https://doi.org/10.1016/s0378-3774(96)01294-2)
- [6] Khan, N., Syeed, S., Masood, A., Nazar, R., and Iqbal, N., Application of salicylic acid increases contents of nutrients and antioxidative metabolism in mungbean and alleviates adverse effects of salinity stress, *Int. J. Plant Sci.*, **1** (2010), e1. <https://doi.org/10.4081/pb.2010.e1>
- [7] Lu, C.M., Qin, N.W., Wang, B.S., Kuang T.Y. Does salt stress lead to increased susceptibility of photosystem II , to photoinhibition and changes in photosynthetic pigment composition in halophyte Suaeda Salsa grown outdoors, *Plant Science*, **163** (2002), 1063-1068. [https://doi.org/10.1016/s0168-9452\(02\)00281-9](https://doi.org/10.1016/s0168-9452(02)00281-9)
- [8] Lutts, S, Kinet, J. M. And Bouharmont, L.,. NaCl-induced senescence in rise (Oryza sativa L.) cultivars differing in salinity resistance, *Annals of Botany*, **78** (1996), 389-398. <https://doi.org/10.1006/anbo.1996.0134>
- [9] Mammadov, G.S, *Basics of Soil Science and Soil Geography*, Baku, Elm, 2007, 664.

- [10] Muhammad Farooq, Mubshar Hussain, Abdul Wakeel, Kadambot H. M. Siddique, Salt stress in maize: effects, resistance mechanisms, and management. A review, *Agronomy for Sustainable Development*, **35** (2015), no. 2, 461-481.
<https://doi.org/10.1007/s13593-015-0287-0>
- [11] Munns R, James RA, A. Läuchli, Approaches to increasing the salt tolerance of wheat and other cereals, *J. Exp. Botany*, **57** (2006), 1025–1043.
<https://doi.org/10.1093/jxb/erj100>
- [12] Perez - Alfocea, F., Balibrea, M.E., Santa Cruz, A. and Estan, M.T. Agronomical and physiological characterization of salinity tolerance in a commercial tomato hybrid, *Plant and Soil.*, **180** (1996), 251-257.
<https://doi.org/10.1007/bf00015308>
- [13] Rustamov, H.N., Talai, J.M., Hasanova, G.M., Ibrahimov, E.R., Ahmadova, G.G. and Musayev, A.J., *Prospects For The Creation Of Intensive Durum Wheat Varieties Under Conditions Of Plain Garabagh*, Collection of scientific works of the Research Institute of Crop Husbandry., 2017, Vol. XXVIII, 86-91.
- [14] Tambussi E. A., Noges S., Araus L. Ear of durum wheat under water stress: Water relations and photosynthetic metabolism, *Planta*, **3** (2005), 446-458.
<https://doi.org/10.1007/s00425-004-1455-7>
- [15] Villora, G., Pulgar, G., Moreno, D.A. and Romero, L., Salinity treatments and their effect on nutrient concentration in zucchini plants (*Cucurbita pepo* L. var. *Moschata*), *Aust. J. Exp. Argi.*, **37** (1997), 605-608.
- [16] Wang Wen – Yuan, Yan Xiao-Feng, Jiang Ying, Qu Bo, Xu Yu –Feng.. Effects of salt stress on water content and photosynthetic characteristics in iris lactera Var. Chinessis seedlings, *Middle – East Journal of Scientific Research*, **12** (2012), no. 1, 70-74.
- [17] Winternmans, J.E.G. and De Mots, A. Spectrophotometric Characteristics of Chlorophyll a and b and Their Phaeophytins in Ethanol, *Biochimica et Biophysica Acta (BBA) - Biophysics including Photosynthesis*, **109** (1965), 448-453.
[https://doi.org/10.1016/0926-6585\(65\)90170-6](https://doi.org/10.1016/0926-6585(65)90170-6)

Received: February 15, 2023; Published: March 9, 2023