

# On the Symmetric Properties for the Generalized Twisted Tangent Polynomials

C. S. Ryoo

Department of Mathematics  
Hannam University, Daejeon 306-791, Korea

Copyright © 2014 C. S. Ryoo. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Abstract

In this paper, we study the symmetry for the generalized twisted tangent numbers  $T_{n,\chi,\zeta}$  and polynomials  $T_{n,\chi,\zeta}(x)$ . Finally, we obtain some interesting identities of the power sums and the generalized twisted polynomials  $T_{n,\chi,\zeta}(x)$  using the symmetric properties for the  $p$ -adic invariant integral on  $\mathbb{Z}_p$ .

**Mathematics Subject Classification:** 11B68, 11S40, 11S80

**Keywords:** The tangent numbers and polynomials, the generalized twisted tangent numbers and polynomials, symmetric properties, power sums

## 1 Introduction

Bernoulli numbers, Bernoulli polynomials, Euler numbers, Euler polynomials, tangent numbers, and tangent polynomials possess many interesting properties and arise in many areas of mathematics and physics(see [1-6]). Throughout this paper we use the following notations. By  $\mathbb{Z}_p$  we denote the ring of  $p$ -adic rational integers,  $\mathbb{Q}$  denotes the field of rational numbers,  $\mathbb{Q}_p$  denotes the field of  $p$ -adic rational numbers,  $\mathbb{C}$  denotes the complex number field, and  $\mathbb{C}_p$  denotes the completion of algebraic closure of  $\mathbb{Q}_p$ . Let  $\nu_p$  be the normalized exponential valuation of  $\mathbb{C}_p$  with  $|p|_p = p^{-\nu_p(p)} = p^{-1}$ . When one talks of  $q$ -extension,  $q$  is considered in many ways such as an indeterminate,

a complex number  $q \in \mathbb{C}$ , or  $p$ -adic number  $q \in \mathbb{C}_p$ . If  $q \in \mathbb{C}$  one normally assume that  $|q| < 1$ . If  $q \in \mathbb{C}_p$ , we normally assume that  $|q - 1|_p < p^{-\frac{1}{p-1}}$  so that  $q^x = \exp(x \log q)$  for  $|x|_p \leq 1$ . Let  $UD(\mathbb{Z}_p)$  be the space of uniformly differentiable function on  $\mathbb{Z}_p$ . For  $g \in UD(\mathbb{Z}_p)$  the fermionic  $p$ -adic invariant  $q$ -integral on  $\mathbb{Z}_p$  is defined by Kim as follows:

$$I_{-q}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-q}(x) = \lim_{N \rightarrow \infty} \frac{1}{[p^N]_{-q}} \sum_{x=0}^{p^N-1} f(x)(-q)^x, \text{ see [2].}$$

Note that

$$\lim_{q \rightarrow 1} I_{-q}(g) = \int_{\mathbb{Z}_p} g(x) d\mu_{-1}(x). \quad (1.1)$$

If we take  $g_n(x) = g(x + n)$  in (1.1), then we see that

$$I_{-1}(g_n) = (-1)^n I_{-1}(g) + 2 \sum_{l=0}^{n-1} (-1)^{n-1-l} g(l). \quad (1.2)$$

Let a fixed positive integer  $d$  with  $(p, d) = 1$ , set

$$X = X_d = \varprojlim_N (\mathbb{Z}/dp^N \mathbb{Z}), \quad X_1 = \mathbb{Z}_p, \quad X^* = \bigcup_{\substack{0 < a < dp \\ (a,p)=1}} a + dp\mathbb{Z}_p,$$

$$a + dp^N \mathbb{Z}_p = \{x \in X \mid x \equiv a \pmod{dp^N}\},$$

where  $a \in \mathbb{Z}$  satisfies the condition  $0 \leq a < dp^N$ . It is easy to see that

$$I_{-1}(g) = \int_X g(x) d\mu_{-1}(x) = \int_{\mathbb{Z}_p} g(x) d\mu_{-1}(x), \text{ see [2].} \quad (1.3)$$

Let  $T_p = \cup_{N \geq 1} C_{p^N} = \lim_{N \rightarrow \infty} C_{p^N}$ , where  $C_{p^N} = \{\zeta \mid \zeta^{p^N} = 1\}$  is the cyclic group of order  $p^N$ . For  $\zeta \in T_p$ , we denote by  $\phi_\zeta : \mathbb{Z}_p \rightarrow \mathbb{C}_p$  the locally constant function  $x \mapsto \zeta^x$ . In [4], we introduced the generalized twisted tangent numbers  $T_{n,\chi,\zeta}$  and polynomials  $T_{n,\chi,\zeta}(x)$  attached to  $\chi$ . Let  $\chi$  be Dirichlet's character with conductor  $d \in \mathbb{N}$  with  $d \equiv 1 \pmod{2}$ . The generalized twisted tangent numbers  $T_{n,\chi,\zeta}$  attached to  $\chi$  and generalized twisted tangent polynomials  $T_{n,\chi,\zeta}(x)$  attached to  $\chi$  are defined by the generating functions

$$\frac{2 \sum_{a=0}^{d-1} \chi(a)(-1)^a \zeta^a e^{2at}}{\zeta^d e^{2dt} + 1} = \sum_{n=0}^{\infty} T_{n,\chi,\zeta} \frac{t^n}{n!}, \quad (1.4)$$

$$\left( \frac{2 \sum_{a=0}^{d-1} \chi(a)(-1)^a \zeta^a e^{2at}}{\zeta^d e^{2dt} + 1} \right) e^{xt} = \sum_{n=0}^{\infty} T_{n,\chi,\zeta}(x) \frac{t^n}{n!}, \text{ cf. [4].} \quad (1.5)$$

**Theorem 1.1** ([4]) For positive integers  $n$  and  $\zeta \in T_p$ , we have

$$T_{n,\chi,\zeta}(x) = \int_X \chi(y) \phi_\zeta(y) (2y+x)^n d\mu_{-1}(y).$$

**Corollary 1.2** ([4]) For positive integers  $n$  and  $\zeta \in T_p$ , we have

$$T_{n,\chi,\zeta} = \int_X \chi(y) \phi_\zeta(y) (2y)^n d\mu_{-1}(y).$$

**Theorem 1.3** ([4]) For positive integers  $n$  and  $\zeta \in T_p$ , we have

$$T_{n,\chi,\zeta}(x) = \sum_{l=0}^n \binom{n}{l} x^{n-l} T_{l,\chi,\zeta}.$$

## 2 Symmetry for the generalized twisted tangent polynomials

In this section, we assume that  $\zeta \in T_p$ . We obtain some interesting identities of the power sums and the generalized twisted polynomials  $T_{n,\chi,\zeta}(x)$  using the symmetric properties for the  $p$ -adic invariant integral on  $\mathbb{Z}_p$ . If  $n$  is odd from (1.2), we obtain

$$I_{-1}(g_n) + I_{-1}(g) = 2 \sum_{k=0}^{n-1} (-1)^k g(k) \quad (\text{see [2]}). \quad (2.1)$$

It will be more convenient to write (2.1) as the equivalent integral form

$$\int_{\mathbb{Z}_p} g(x+n) d\mu_{-1}(x) + \int_{\mathbb{Z}_p} g(x) d\mu_{-1}(x) = 2 \sum_{k=0}^{n-1} (-1)^k g(k). \quad (2.2)$$

Substituting  $g(x) = \chi(x) \zeta^x e^{2xt}$  into the above, we obtain

$$\begin{aligned} & \int_X \chi(x+n) \zeta^{x+n} e^{(2x+2n)t} d\mu_{-1}(x) + \int_X \chi(x) \zeta^x e^{2xt} d\mu_{-1}(x) \\ &= 2 \sum_{j=0}^{n-1} (-1)^j \chi(j) \zeta^j e^{2jt}. \end{aligned} \quad (2.3)$$

Let us define the alternating power sums  $\mathcal{T}_{k,\chi,\zeta}(n)$  as follows:

$$\mathcal{T}_{k,\chi,\zeta}(n) = \sum_{l=0}^n (-1)^l \chi(l) \zeta^l (2l)^k. \quad (2.4)$$

After some elementary calculations, we have

$$\begin{aligned} & \int_X \chi(x) \zeta^{x+nd} e^{(2x+2nd)t} d\mu_{-1}(x) + \int_X \chi(x) \zeta^x e^{2xt} d\mu_{-1}(x) \\ &= (1 + \zeta^{nd} e^{2ndt}) \frac{2 \sum_{a=0}^{d-1} \chi(a) (-1)^a \zeta^a e^{2at}}{\zeta^d e^{2dt} + 1}. \end{aligned} \quad (2.5)$$

From the above, we get

$$\begin{aligned} & \int_X \chi(x) \zeta^{x+nd} e^{(2x+2nd)t} d\mu_{-1}(x) + \int_X \chi(x) \zeta^x e^{2xt} d\mu_{-1}(x) \\ &= \frac{2 \int_X \chi(x) \zeta^x e^{2xt} d\mu_{-1}(x)}{\int_X \zeta^{ndx} e^{2ndtx} d\mu_{-1}(x)}. \end{aligned} \quad (2.6)$$

By substituting Taylor series of  $e^{2xt}$  into (2.3) and by using (2.4), we have

$$\begin{aligned} & \zeta^{nd} \sum_{k=0}^m \binom{m}{k} (2nd)^{m-k} \int_X \chi(x) \zeta^x (2x)^k d\mu_{-1}(x) + \int_X \chi(x) \zeta^x (2x)^m d\mu_{-1}(x) \\ &= 2\mathcal{T}_{m,\chi,\zeta}(nd-1). \end{aligned} \quad (2.7)$$

By using (2.6) and (2.7), we arrive at the following theorem:

**Theorem 2.1** *Let  $n$  be odd positive integer. Then we obtain*

$$\frac{2 \int_X \chi(x) \zeta^x e^{2xt} d\mu_{-1}(x)}{\int_X \zeta^{ndx} e^{2ndtx} d\mu_{-1}(x)} = \sum_{m=0}^{\infty} (2\mathcal{T}_{m,\chi,\zeta}(nd-1)) \frac{t^m}{m!}.$$

Let  $w_1$  and  $w_2$  be odd positive integers. Then we set

$$\begin{aligned} S(w_1, w_2) &= \\ &= \frac{\int_X \int_X \chi(x_1) \chi(x_2) \zeta^{(w_1 x_1 + w_2 x_2)} e^{(2w_1 x_1 + 2w_2 x_2 + w_1 w_2 x) t} d\mu_{-1}(x_1) d\mu_{-1}(x_2)}{\int_X \zeta^{w_1 w_2 dx} e^{2w_1 w_2 dxt} d\mu_{-1}(x)}. \end{aligned} \quad (2.8)$$

By Theorem 2.1 and (2.8), after elementary calculations, we obtain

$$S(w_1, w_2) = \left( \frac{1}{2} \sum_{m=0}^{\infty} T_{m,\chi,\zeta^{w_1}}(w_2 x) w_1^m \frac{t^m}{m!} \right) \left( 2 \sum_{m=0}^{\infty} \mathcal{T}_{m,\chi,\zeta^{w_2}}(w_1 d - 1) w_2^m \frac{t^m}{m!} \right).$$

By using Cauchy product in the above, we have

$$S(w_1, w_2) = \sum_{m=0}^{\infty} \left( \sum_{j=0}^m \binom{m}{j} T_{j,\chi,\zeta^{w_1}}(w_2 x) w_1^j \mathcal{T}_{m-j,\chi,\zeta^{w_2}}(w_1 d - 1) w_2^{m-j} \right) \frac{t^m}{m!}. \quad (2.9)$$

From the symmetry of  $S(w_1, w_2)$  in  $w_1$  and  $w_2$ , we also see that

$$S(w_1, w_2) = \left( \frac{1}{2} \sum_{m=0}^{\infty} T_{m, \chi, \zeta^{w_2}}(w_1 x) w_2^m \frac{t^m}{m!} \right) \left( 2 \sum_{m=0}^{\infty} \mathcal{T}_{m, \chi, \zeta^{w_1}}(w_2 d - 1) w_1^m \frac{t^m}{m!} \right).$$

Thus we have

$$S(w_1, w_2) = \sum_{m=0}^{\infty} \left( \sum_{j=0}^m \binom{m}{j} T_{j, \chi, \zeta^{w_2}}(w_1 x) w_2^j \mathcal{T}_{m-j, \chi, \zeta^{w_1}}(w_2 d - 1) w_1^{m-j} \right) \frac{t^m}{m!} \quad (2.10)$$

By comparing coefficients  $\frac{t^m}{m!}$  in the both sides of (2.9) and (2.10), we arrive at the following theorem:

**Theorem 2.2** *Let  $w_1$  and  $w_2$  be odd positive integers. Then we obtain*

$$\begin{aligned} & \sum_{j=0}^m \binom{m}{j} w_1^{m-j} w_2^j T_{j, \chi, \zeta^{w_2}}(w_1 x) \mathcal{T}_{m-j, \chi, \zeta^{w_1}}(w_2 d - 1) \\ &= \sum_{j=0}^m \binom{m}{j} w_1^j w_2^{m-j} T_{j, \chi, \zeta^{w_1}}(w_2 x) \mathcal{T}_{m-j, \chi, \zeta^{w_2}}(w_1 d - 1), \end{aligned}$$

where  $T_{k, \chi, \zeta}(x)$  and  $\mathcal{T}_{m, \chi, \zeta}(k)$  denote the generalized twisted tangent polynomials and the alternating sums of powers, respectively.

By Theorem 2.2 and Theorem 1.3, we have the following corollary.

**Corollary 2.3** *Let  $w_1$  and  $w_2$  be odd positive integers. Then we obtain*

$$\begin{aligned} & \sum_{j=0}^m \sum_{k=0}^j \binom{m}{j} \binom{j}{k} w_1^{m-k} w_2^j x^{j-k} T_{k, \chi, \zeta^{w_2}} \mathcal{T}_{m-j, \chi, \zeta^{w_1}}(w_2 d - 1) \\ &= \sum_{j=0}^m \sum_{k=0}^j \binom{m}{j} \binom{j}{k} w_1^j w_2^{m-k} x^{j-k} T_{k, \chi, \zeta^{w_1}} \mathcal{T}_{m-j, \chi, \zeta^{w_2}}(w_1 d - 1). \end{aligned}$$

Now we will derive another interesting identities for the generalized twisted tangent polynomials using the symmetric property of  $S(w_1, w_2)$ . By (2.8),

after elementary calculations, we have

$$\begin{aligned}
 S(w_1, w_2) &= \left( \frac{1}{2} e^{w_1 w_2 x t} \int_X \chi(x_1) \zeta^{w_1 x_1} e^{2x_1 w_1 t} d\mu_{-1}(x_1) \right) \left( 2 \sum_{j=0}^{w_1 d-1} (-1)^j \chi(j) \zeta^{w_2 j} e^{2j w_2 t} \right) \\
 &= \sum_{j=0}^{w_1 d-1} (-1)^j \chi(j) \zeta^{w_2 j} \int_X \chi(x_1) \zeta^{w_1 x_1} e^{\left( 2x_1 + w_2 x + \frac{2j w_2}{w_1} \right) (w_1 t)} d\mu_{-1}(x_1) \\
 &= \sum_{n=0}^{\infty} \left( \sum_{j=0}^{w_1 d-1} (-1)^j \chi(j) \zeta^{w_2 j} T_{n, \chi, \zeta^{w_1}} \left( w_2 x + \frac{2j w_2}{w_1} \right) w_1^n \right) \frac{t^n}{n!}.
 \end{aligned} \tag{2.11}$$

By using the symmetry property in (2.11), we also have

$$\begin{aligned}
 S(w_1, w_2) &= \sum_{j=0}^{w_2 d-1} (-1)^j \chi(j) \zeta^{w_1 j} \int_X \chi(x_2) \zeta^{w_2 x_2} e^{\left( 2x_2 + w_1 x + \frac{2j w_1}{w_2} \right) (w_2 t)} d\mu_{-1}(x_2) \\
 &= \sum_{n=0}^{\infty} \left( \sum_{j=0}^{w_2 d-1} (-1)^j \chi(j) \zeta^{w_1 j} T_{n, \chi, \zeta^{w_2}} \left( w_1 x + \frac{2j w_1}{w_2} \right) w_2^n \right) \frac{t^n}{n!}.
 \end{aligned} \tag{2.12}$$

By comparing coefficients  $\frac{t^n}{n!}$  in the both sides of (2.11) and (2.12), we have the following theorem.

**Theorem 2.4** *Let  $w_1$  and  $w_2$  be odd positive integers. Then we obtain*

$$\begin{aligned}
 &\sum_{j=0}^{w_1 d-1} (-1)^j \chi(j) \zeta^{w_2 j} T_{n, \chi, \zeta^{w_1}} \left( w_2 x + \frac{2j w_2}{w_1} \right) w_1^n \\
 &= \sum_{j=0}^{w_2 d-1} (-1)^j \chi(j) \zeta^{w_1 j} T_{n, \chi, \zeta^{w_2}} \left( w_1 x + \frac{2j w_1}{w_2} \right) w_2^n.
 \end{aligned}$$

If we take  $x = 0$  in Theorem 2.4, we also derive the interesting identity for the generalized twisted tangent numbers as follows:

$$\begin{aligned}
 &\sum_{j=0}^{w_1 d-1} (-1)^j \chi(j) \zeta^{w_2 j} T_{n, \chi, \zeta^{w_1}} \left( \frac{2w_2 j}{w_1} \right) w_1^n \\
 &= \sum_{j=0}^{w_2 d-1} (-1)^j \chi(j) \zeta^{w_1 j} T_{n, \chi, \zeta^{w_2}} \left( \frac{2w_1 j}{w_2} \right) w_2^n.
 \end{aligned}$$

Letting  $\zeta \rightarrow 1$  in Theorem 2.4, we can immediately have the generalized multiplication theorem for the generalized tangent polynomials(see, [5]).

## References

- [1] L. Comtet, *Advances combinatorics*, Riedel, Dordrecht, 1974.
- [2] T. Kim, *q-Volkenborn integration*, Russ. J. Math. Phys., **9**(2002), 288-299.
- [3] C. S. Ryoo, *A note on the tangent numbers and polynomials*, Adv. Studies Theor. Phys., **7(9)**(2013), 447 - 454.
- [4] C. S. Ryoo, *A Note on the Generalized Twisted Tangent Polynomials*, Int. Journal of Math. Analysis, **7(55)**(2013), 2717 - 2722.
- [5] C. S. Ryoo, *Symmetric Identities for the Generalized Tangent Polynomials Associated with p-Adic Integral on  $\mathbb{Z}_p$* , Applied Mathematical Sciences, **8(17)**(2014), 829-835.
- [6] H. Shin, J. Zeng, *The q-tangent and q-secant numbers via continued fractions*, European J. Combin., **31**(2010), 1689-1705.

**Received: September 15, 2014; Published: October 29, 2014**