

# Hankel Determinant for Starlike and Convex Functions

**Aini Janteng**

School of Science and Technology, Universiti Malaysia Sabah  
Locked Bag No.2073, 88999 Kota Kinabalu, Sabah, Malaysia  
aini\_jg@ums.edu.my

**Suzeini Abdul Halim**

Institute of Mathematical Sciences, Universiti Malaya  
50603 Kuala Lumpur, Malaysia  
suzeini@um.edu.my

**Maslina Darus**

School of Mathematical Sciences  
Faculty of Sciences and Technology  
Universiti Kebangsaan Malaysia  
43600 Bangi, Selangor, Malaysia  
maslina@pkrisc.cc.ukm.my

## Abstract

Denote  $\mathcal{S}$  to be the class of functions which are analytic, normalised and univalent in the open unit disc  $\mathcal{D} = \{z : |z| < 1\}$ . The important subclasses of  $\mathcal{S}$  are the class of starlike and convex functions, which we denote by  $\mathcal{S}^*$  and  $\mathcal{C}$ . This paper focuses on attaining sharp upper bounds for the functional  $|a_2a_4 - a_3^2|$  for functions  $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$  belonging to  $\mathcal{S}^*$  and  $\mathcal{C}$ .

**Mathematics Subject Classification:** Primary 30C45

**Keywords:** starlike functions, convex functions, Hankel determinant

# 1 Introduction

Let  $\mathcal{S}$  denote the class of normalised analytic univalent functions  $f$  of the form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n \quad (1)$$

where  $z \in \mathcal{D} = \{z : |z| < 1\}$ . In [5], the  $q$ th Hankel determinant for  $q \geq 1$  and  $n \geq 0$  is stated by Noonan and Thomas as

$$H_q(n) = \begin{vmatrix} a_n & a_{n+1} \dots & a_{n+q+1} \\ a_{n+1} & \dots & \vdots \\ \vdots & & \\ a_{n+q-1} & \dots & a_{n+2q-2} \end{vmatrix}.$$

This determinant has also been considered by several authors. For example, Noor in [6] determined the rate of growth of  $H_q(n)$  as  $n \rightarrow \infty$  for functions  $f$  given by (1) with bounded boundary. Ehrenborg in [1] studied the Hankel determinant of exponential polynomials. The Hankel transform of an integer sequence and some of its properties were discussed by Layman in [4].

Easily, one can observe that the Fekete and Szegő functional is  $\mathcal{H}_2(1)$ . Fekete and Szegő then further generalised the estimate  $|a_3 - \mu a_2^2|$  where  $\mu$  is real and  $f \in \mathcal{S}$ . For our discussion in this paper, we consider the Hankel determinant in the case  $q = 2$  and  $n = 2$ ,

$$H_2(2) = \begin{vmatrix} a_2 & a_3 \\ a_3 & a_4 \end{vmatrix}.$$

We seek upper bound for the functional  $|a_2 a_4 - a_3^2|$  for functions  $f$  belongs to the class  $\mathcal{S}^*$  and  $\mathcal{C}$ . The class  $\mathcal{S}^*$  and  $\mathcal{C}$  are defined as follows.

**Definition 1.1** *Let  $f$  be given by (1). Then  $f \in \mathcal{S}^*$  if and only if*

$$\operatorname{Re} \left\{ \frac{z f'(z)}{f(z)} \right\} > 0, \quad z \in \mathcal{D}. \quad (2)$$

**Definition 1.2** *Let  $f$  be given by (1). Then  $f \in \mathcal{C}$  if and only if*

$$\operatorname{Re} \left\{ \frac{(z f'(z))'}{f'(z)} \right\} > 0, \quad z \in \mathcal{D}. \quad (3)$$

It follows that  $f \in \mathcal{C}$  if and only if  $z f'(z) \in \mathcal{S}^*$ .

First, some preliminary lemmas.

## 2 Preliminary Results

Let  $\mathcal{P}$  be the family of all functions  $p$  analytic in  $\mathcal{D}$  for which  $Re\{p(z)\} > 0$  and

$$p(z) = 1 + c_1z + c_2z^2 + \dots \tag{4}$$

for  $z \in \mathcal{D}$ .

**Lemma 2.1** ([7]) *If  $p \in \mathcal{P}$  then  $|c_k| \leq 2$  for each  $k$ .*

**Lemma 2.2** ([3]) *The power series for  $p$  given in (4) converges in  $\mathcal{D}$  to a function in  $\mathcal{P}$  if and only if the Toeplitz determinants*

$$D_n = \begin{vmatrix} 2 & c_1 & c_2 \dots & c_n \\ c_{-1} & 2 & c_1 \dots & c_{n-1} \\ \vdots & & & \\ c_{-n} & c_{-n+1} & c_{-n+2} \dots & 2 \end{vmatrix}, \quad n = 1, 2, 3, \dots \tag{5}$$

and  $c_{-k} = \bar{c}_k$ , are all nonnegative. They are strictly positive except for  $p(z) = \sum_{k=1}^m \rho_k p_o(e^{it_k} z)$ ,  $\rho_k > 0$ ,  $t_k$  real and  $t_k \neq t_j$  for  $k \neq j$ ; in this case  $D_n > 0$  for  $n < m - 1$  and  $D_n = 0$  for  $n \geq m$ .

This necessary and sufficient condition is due to Carathéodory and Toeplitz and can be found in [3].

## 3 Main Result

**Theorem 3.1** *Let  $f \in \mathcal{S}^*$ . Then*

$$|a_2a_4 - a_3^2| \leq 1.$$

*The result obtained is sharp.*

**Proof.**

Since  $f \in \mathcal{S}^*$ , it follows from (2) that  $\exists p \in \mathcal{P}$  such that

$$zf'(z) = f(z)p(z) \tag{6}$$

for some  $z \in \mathcal{D}$ . Equating coefficients in (6) yields

$$\left. \begin{aligned} a_2 &= c_1 \\ a_3 &= \frac{c_2}{2} + \frac{c_1^2}{2} \\ a_4 &= \frac{c_3}{3} + \frac{c_1c_2}{2} + \frac{c_1^3}{6} \end{aligned} \right\}. \tag{7}$$

From (7), it is easily established that

$$|a_2a_4 - a_3^2| = \left| \frac{c_1c_3}{3} - \frac{c_2^2}{4} - \frac{c_1^4}{12} \right|. \quad (8)$$

Lemma 2.2 can then be used to obtain the proper bound on (8). We may assume without restriction that  $c_1 \geq 0$ . Rewriting (5) for the cases  $n=2$  and  $n=3$ , result in

$$D_2 = \begin{vmatrix} 2 & c_1 & c_2 \\ c_1 & 2 & c_1 \\ \bar{c}_2 & c_1 & 2 \end{vmatrix} = 8 + 2 \operatorname{Re}\{c_1^2c_2\} - 2|c_2|^2 - 4c_1^2 \geq 0,$$

which is equivalent to

$$2c_2 = c_1^2 + x(4 - c_1^2) \quad (9)$$

for some  $x, |x| \leq 1$ .

Further,  $D_3 \geq 0$  is equivalent to

$$|(4c_3 - 4c_1c_2 + c_1^3)(4 - c_1^2) + c_1(2c_2 - c_1^2)^2| \leq 2(4 - c_1^2)^2 - 2|2c_2 - c_1^2|^2;$$

and this, with (9), provides the relation

$$4c_3 = c_1^3 + 2(4 - c_1^2)c_1x - c_1(4 - c_1^2)x^2 + 2(4 - c_1^2)(1 - |x|^2)z, \quad (10)$$

for some value of  $z, |z| \leq 1$ .

Suppose now that  $c_1 = c$  and  $0 \leq c \leq 2$ . Using (9) along with (10), we obtain

$$\begin{aligned} \left| \frac{c_1c_3}{3} - \frac{c_2^2}{4} - \frac{c_1^4}{12} \right| &= \left| \frac{(4 - c^2)c^2x}{24} - \frac{c^4}{16} \right. \\ &\quad \left. + \frac{(4 - c^2)(1 - |x|^2)cz}{6} - \frac{(4 - c^2)x^2(12 + c^2)}{48} \right|. \end{aligned}$$

Application of the triangle inequality gives

$$\begin{aligned} \left| \frac{c_1c_3}{3} - \frac{c_2^2}{4} - \frac{c_1^4}{12} \right| &\leq \frac{c^4}{16} + \frac{c(4 - c^2)}{6} + \frac{c^2(4 - c^2)\rho}{24} \\ &\quad + \frac{(4 - c^2)(c - 2)(c - 6)\rho^2}{48} \\ &= F(\rho) \end{aligned} \quad (11)$$

with  $\rho = |x| \leq 1$ . Furthermore,

$$F'(\rho) = \frac{c^2(4 - c^2)}{24} + \frac{(4 - c^2)(c - 2)(c - 6)\rho}{24}$$

and with elementary calculus, one can show that  $F'(\rho) > 0$  for  $\rho > 0$ ; implying that  $F$  is an increasing function and thus the upper bound for (11) corresponds to  $\rho = 1$ , in which case

$$\left| \frac{c_1 c_3}{3} - \frac{c_2^2}{4} - \frac{c_1^4}{12} \right| \leq 1$$

for all  $c \in [0, 2]$ . Equality is attained for functions in  $\mathcal{S}^*$  given by

$$\frac{zf'(z)}{f(z)} = \frac{1+z}{1-z}$$

and

$$\frac{zf'(z)}{f(z)} = \frac{1+z^2}{1-z^2}.$$

This completes the proof of theorem 3.1.

**Theorem 3.2** *Let  $f \in \mathcal{C}$ . Then*

$$|a_2 a_4 - a_3^2| \leq \frac{1}{8}.$$

*The result obtained is sharp.*

**Proof.**

Similar approach as in the proof of Theorem 3.1. Since  $f \in \mathcal{C}$ , it follows from (3) that  $\exists p \in \mathcal{P}$  such that

$$(zf'(z))' = f'(z)p(z) \tag{12}$$

for some  $z \in \mathcal{D}$ . Equating coefficients in (12) yields

$$\left. \begin{aligned} a_2 &= \frac{c_1}{2} \\ a_3 &= \frac{c_2}{6} + \frac{c_1^2}{6} \\ a_4 &= \frac{c_3}{12} + \frac{c_1 c_2}{8} + \frac{c_1^3}{24} \end{aligned} \right\}. \tag{13}$$

From (13), it is easily established that

$$|a_2 a_4 - a_3^2| = \frac{1}{144} |6c_1 c_3 + c_1^2 c_2 - 4c_2^2 - c_1^4|. \tag{14}$$

Now, assuming  $c_1 = c(0 \leq c \leq 2)$  and using (9) together with (10) we have

$$\left| 6c_1 c_3 + c_1^2 c_2 - 4c_2^2 - c_1^4 \right| = \left| \frac{3c^2(4-c^2)x}{2} - \frac{(4-c^2)(8+c^2)x^2}{2} + 3c(4-c^2)(1-|x|^2)z \right|$$

and an application of the triangle inequality shows that

$$\begin{aligned} \left| 6c_1c_3 + c_1^2c_2 - 4c_2^2 - c_1^4 \right| &\leq 3c(4 - c^2) + \frac{3c^2(4 - c^2)\rho}{2} \\ &\quad + \frac{(4 - c^2)(c - 2)(c - 4)\rho^2}{2} \\ &= F(\rho) \end{aligned} \tag{15}$$

with  $\rho = |x| \leq 1$ . For

$$F'(\rho) = \frac{3c^2(4 - c^2)}{2} + (c - 2)(c - 4)(4 - c^2)\rho,$$

it can be shown that  $F'(\rho) > 0$  and thus is an increasing function implying  $\text{Max}_{\rho \leq 1} F(\rho) = F(1)$ . Now let

$$\begin{aligned} G(c) &= F(1) \\ &= 3c(4 - c^2) + \frac{3c^2(4 - c^2)}{2} + \frac{(4 - c^2)(c - 2)(c - 4)}{2}. \end{aligned}$$

Trivially, one can show that  $G$  has a maximum attained at  $c = 1$ . The upper bound for (15) corresponds to  $\rho = 1$  and  $c = 1$ , in which case

$$\left| 6c_1c_3 + c_1^2c_2 - 4c_2^2 - c_1^4 \right| \leq 18.$$

Letting  $c_1 = 1$ ,  $c_2 = -1$  and  $c_3 = -2$  in (14) shows that the result is sharp. This completes the proof of Theorem 3.2.

## References

- [1] Ehrenborg, R. : The Hankel determinant of exponential polynomials, *American Mathematical Monthly*, **107**(2000): 557-560.
- [2] Fekete, M. and Szegő, G. : Eine Bemerkung uber ungerade schlichte Funktionen, *J. London Math. Soc.*, **8** (1933): 85-89.
- [3] Grenander, U. and Szegő, G. : Toeplitz forms and their application, *Univ. of California Press, Berkeley and Los Angeles*, (1958)
- [4] Layman, J.W. : The Hankel transform and some of its properties, *J. of integer sequences*, **4**(2001): 1-11.

- [5] Noonan, J.W. and Thomas, D.K. : On the second Hankel determinant of areally mean  $p$ -valent functions, *Trans. Amer. Math. Soc.*, **223** (2)(1976): 337-346.
- [6] Noor, K.I. : Hankel determinant problem for the class of functions with bounded boundary rotation, *Rev. Roum. Math. Pures Et Appl.*, **28**(8)(1983): 731-739.
- [7] Pommerenke, Ch. : Univalent functions, *Vandenhoeck and Ruprecht, Göttingen*, (1975)

**Received: January 20, 2007**